

ASYMPTOTIC STABILITY AND DECAY RATES OF HOMOGENEOUS POSITIVE SYSTEMS WITH BOUNDED AND UNBOUNDED DELAYS*

HAMID REZA FEYZMAHDAVIAN[†], THEMISTOKLIS CHARALAMBOUS[†], AND
MIKAEL JOHANSSON[†]

Abstract. There are several results on the stability of nonlinear positive systems in the presence of time delays. However, most of them assume that the delays are constant. This paper considers *time-varying*, possibly unbounded, delays and establishes asymptotic stability and bounds the decay rate of a significant class of nonlinear positive systems which includes positive linear systems as a special case. Specifically, we present a necessary and sufficient condition for delay-independent stability of continuous-time positive systems whose vector fields are cooperative and homogeneous. We show that global asymptotic stability of such systems is independent of the magnitude and variation of the time delays. For various classes of time delays, we are able to derive explicit expressions that quantify the decay rates of positive systems. We also provide the corresponding counterparts for discrete-time positive systems whose vector fields are nondecreasing and homogeneous.

Key words. monotone system, positive system, homogeneous system, time-varying delay

AMS subject classifications. 34C12, 34K20, 93C23, 93D05

DOI. 10.1137/130943340

1. Introduction. Many real-world processes in areas such as economics, biology, ecology, and communications deal with physical quantities that cannot attain negative values. The state trajectories of dynamical models characterizing such processes should thus be constrained to stay within the positive orthant. Such systems are commonly referred to as *positive systems* [41, 11, 18]. Due to their importance and broad applications, a large body of literature has been concerned with the analysis and control of positive systems (see, e.g., [22, 44, 45, 24, 1, 37, 30, 34, 10, 23, 38, 43, 14, 25, 5] and references therein).

In distributed systems where exchange of information is involved, delays are inevitable. For this reason, a considerable effort has been devoted to characterizing the stability and performance of systems with delays (see, e.g., [19, 15, 46, 16, 20, 35, 40] and references therein). Recently, the stability of delayed positive linear systems has received significant attention [17, 32, 28, 36, 29] and it has been shown that such systems are insensitive to certain classes of time delays, in the sense that a positive linear system with time delays is asymptotically stable if the corresponding delay-free system is asymptotically stable. This is a surprising property, since the stability of general dynamical systems typically depends on the magnitude and variation of the time delays.

While the asymptotic stability of positive linear systems in the presence of time delays has been thoroughly investigated, the theory for *nonlinear* positive systems is considerably less well-developed (see, e.g., [17] and [31, 4] for exceptions). In particu-

*Received by the editors October 29, 2013; accepted for publication (in revised form) June 20, 2014; published electronically August 28, 2014.

<http://www.siam.org/journals/sicon/52-4/94334.html>

[†]Department of Automatic Control, School of Electrical Engineering and ACCESS Linnaeus Center, Royal Institute of Technology (KTH), SE-100 44 Stockholm, Sweden (hamidrez@kth.se, themisc@kth.se, mikaelj@kth.se). This work was supported in part by the Swedish Foundation for Strategic Research (SSF) and the Swedish Science Foundation (VR).

lar, [31] showed that the asymptotic stability of a particular class of nonlinear positive systems whose vector fields are *cooperative* and *homogenous* of degree zero does not depend on the magnitude of *constant* delays. A similar result for cooperative systems that are homogeneous of any degree was given in [4], also under the assumption of constant delays. Extensions of these results to *time-varying* delays are, however, not trivial. The main reason for this is that the proof technique in [31, 4] relies on a fundamental monotonicity property of trajectories of cooperative systems, which does not hold when the delays are time-varying. To the best of our knowledge, there have been rather few studies on stability of nonlinear positive systems with time-varying delays (see, e.g., [33, 13]).

At this point, it is worth noting that the results for positive linear systems cited above consider *bounded* delays. However, in some cases, it is not possible to a priori guarantee that the delays will be bounded, but the state evolution might be affected by the entire history of states. It is then natural to ask if the insensitivity properties of positive linear systems with respect to time delays will hold also for *unbounded* delays. In [26], it was shown that, for a particular class of unbounded delays, this is indeed the case. Extensions of this result to more general classes of unbounded delays were given in [42, 12] for continuous- and discrete-time positive linear systems, respectively. However, [26, 42, 12] did not quantify how various bounds on the delay evolution impact the decay rate of positive linear systems.

This paper establishes delay-independent stability of a class of nonlinear positive systems, which includes positive linear systems as a special case, and allows for time-varying, possibly unbounded, delays. The proof technique, which uses neither the Lyapunov–Krasovskii functional method widely used to analyze positive systems with constant delays [17] nor the approach used in [31, 4], allows us to impose minimal restrictions on the delays. Specifically, we make the following contributions:

1. We derive a set of necessary and sufficient conditions for delay-independent global stability of (i) continuous-time positive systems whose vector fields are cooperative and homogeneous of *arbitrary degree* and (ii) discrete-time positive systems whose vector fields are nondecreasing and homogeneous of *degree zero*. We demonstrate that such systems are insensitive to a general class of time delays which includes bounded and unbounded time-varying delays.
2. When the asymptotic behavior of the time delays is known, we obtain conditions to ensure global μ -stability in the sense of [7]. These results allow us to quantify the decay rates of positive systems for various classes of (possibly unbounded) time-varying delays.
3. For bounded delays and a particular class of unbounded delays, we present explicit bounds on the decay rates. These bounds quantify how the magnitude of bounded delays and the rate at which the unbounded delays grow large affect the decay rate.
4. We also show that discrete-time positive systems whose vector fields are nondecreasing and homogeneous of *degree greater than zero* are locally asymptotically stable under delay-independent global stability conditions that we have derived.

The remainder of the paper is organized as follows. In section 2, we introduce the notation and review some preliminaries that are essential for the development of the results in this paper. Our main results for continuous- and discrete-time nonlinear positive systems are stated in sections 3 and 4, respectively. An illustrative example, justifying the validity of our results, is presented in section 5. Finally, concluding remarks are given in section 6.

2. Notation and preliminaries.

2.1. Notation. Vectors are written in bold lower case letters and matrices in capital letters. We let \mathbb{R} , \mathbb{N} , and \mathbb{N}_0 denote the set of real numbers, natural numbers, and the set of natural numbers including zero, respectively. The nonnegative orthant of the n -dimensional real space \mathbb{R}^n is represented by \mathbb{R}_+^n . The i th component of a vector $\mathbf{x} \in \mathbb{R}^n$ is denoted by x_i , and the notation $\mathbf{x} \geq \mathbf{y}$ means that $x_i \geq y_i$ for all components i . If \mathbf{v} is a vector in \mathbb{R}^n , the notation $\mathbf{v} > \mathbf{0}$ indicates that all components of \mathbf{v} are positive. Given a vector $\mathbf{v} > \mathbf{0}$, the weighted l_∞ norm is defined by

$$\|\mathbf{x}\|_\infty^{\mathbf{v}} = \max_{1 \leq i \leq n} \frac{|x_i|}{v_i}.$$

For a matrix $A \in \mathbb{R}^{n \times n}$, a_{ij} denotes the real-valued entry in row i and column j . A matrix $A \in \mathbb{R}^{n \times n}$ is said to be *nonnegative* if $a_{ij} \geq 0$ for all i and j . It is called *Metzler* if $a_{ij} \geq 0$ for all $i \neq j$. Given an n -tuple $\mathbf{r} = (r_1, \dots, r_n)$ of positive real numbers and $\lambda > 0$, the *dilation map* $\delta_\lambda^{\mathbf{r}}(\mathbf{x}) : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is given by

$$\delta_\lambda^{\mathbf{r}}(\mathbf{x}) = (\lambda^{r_1}x_1, \dots, \lambda^{r_n}x_n).$$

If $\mathbf{r} = (1, \dots, 1)$, the dilation map is called the *standard dilation map*. For a real interval $[a, b]$, $\mathcal{C}([a, b], \mathbb{R}^n)$ denotes the space of all real-valued continuous functions on $[a, b]$ taking values in \mathbb{R}^n . The upper-right Dini-derivative of a continuous function $h : \mathbb{R} \rightarrow \mathbb{R}$ at $t = t_0$ is defined by

$$D^+h(t)|_{t=t_0} = \lim_{\Delta \rightarrow 0^+} \sup \frac{h(t_0 + \Delta) - h(t_0)}{\Delta},$$

where $\Delta \rightarrow 0^+$ means that Δ approaches zero from the right-hand side.

2.2. Preliminaries. Next, we review the key definitions and results necessary for developing the main results of this paper. We start with the definition of *cooperative* vector fields.

DEFINITION 2.1. A continuous vector field $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ which is continuously differentiable on $\mathbb{R}^n \setminus \{\mathbf{0}\}$ is said to be *cooperative* if the Jacobian matrix $\partial f / \partial \mathbf{x}$ is Metzler for all $\mathbf{x} \in \mathbb{R}_+^n \setminus \{\mathbf{0}\}$.

Cooperative vector fields satisfy the following property.

PROPOSITION 2.2 (see [41, Remark 3.1.1]). Let $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be cooperative. For any two vectors \mathbf{x} and \mathbf{y} in $\mathbb{R}_+^n \setminus \{\mathbf{0}\}$ with $x_i = y_i$ and $\mathbf{x} \geq \mathbf{y}$, we have $f_i(\mathbf{x}) \geq f_i(\mathbf{y})$.

The following definition introduces *homogeneous* vector fields.

DEFINITION 2.3. For any $p \geq 0$, the vector field $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is said to be *homogeneous of degree p with respect to the dilation map $\delta_\lambda^{\mathbf{r}}(\mathbf{x})$* if

$$\mathbf{f}(\delta_\lambda^{\mathbf{r}}(\mathbf{x})) = \lambda^p \delta_\lambda^{\mathbf{r}}(\mathbf{f}(\mathbf{x})) \quad \forall \mathbf{x} \in \mathbb{R}^n, \forall \lambda > 0.$$

Finally, we define *nondecreasing* vector fields.

DEFINITION 2.4. A vector field $g : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is said to be *nondecreasing on \mathbb{R}_+^n* if $g(\mathbf{x}) \geq g(\mathbf{y})$ for any $\mathbf{x}, \mathbf{y} \in \mathbb{R}_+^n$ such that $\mathbf{x} \geq \mathbf{y}$.

3. Continuous-time homogeneous cooperative systems.

3.1. Problem statement. Consider the continuous-time dynamical system

$$(3.1) \quad \mathcal{G} : \begin{cases} \dot{\mathbf{x}}(t) = \mathbf{f}(\mathbf{x}(t)) + \mathbf{g}(\mathbf{x}(t - \tau(t))), & t \geq 0, \\ \mathbf{x}(t) = \boldsymbol{\varphi}(t), & t \in [-\tau_{\max}, 0], \end{cases}$$

where $\mathbf{x}(t) \in \mathbb{R}^n$ is the state variable, and $f, g : \mathbb{R}^n \rightarrow \mathbb{R}^n$ are continuous vector fields on \mathbb{R}^n , continuously differentiable on $\mathbb{R}^n \setminus \{\mathbf{0}\}$, and such that $\mathbf{f}(\mathbf{0}) = \mathbf{g}(\mathbf{0}) = \mathbf{0}$. Here, $\tau_{\max} \geq 0$, $\varphi(\cdot) \in \mathcal{C}([-\tau_{\max}, 0], \mathbb{R}^n)$ is the vector-valued function specifying the initial condition of the system, and $\tau(\cdot)$ is the time-varying delay which satisfies the following assumption.

Assumption 3.1. The delay $\tau : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is continuous with respect to time and satisfies

$$(3.2) \quad \lim_{t \rightarrow +\infty} t - \tau(t) = +\infty.$$

Note that $\tau(t)$ is not necessarily continuously differentiable and no restriction on its derivative (such as $\dot{\tau}(t) < 1$) is imposed. Condition (3.2) implies that as t increases, $\tau(t)$ grows slower than time itself. This constraint on time delays is not restrictive and typically satisfied in real-world applications. For example, the continuous-time power control algorithm for a wireless network consisting of n mobile users can be described by

$$(3.3) \quad \dot{x}_i(t) = -x_i(t) + \sum_{\substack{j=1 \\ j \neq i}}^n a_{ij} x_j(t - \tau(t)), \quad i = 1, \dots, n.$$

Here, x_i represents the transmit power of user i , and a_{ij} are nonnegative constants [6, 47]. If $\tau(t)$ satisfies (3.2), then given any time $t_1 \geq 0$, there exists a time $t_2 > t_1$ such that

$$t - \tau(t) \geq t_1 \quad \forall t \geq t_2.$$

This simply means that given any time t_1 , information about which transmit power each user has applied prior to t_1 will be received by every other user before a sufficiently long time t_2 and not be used in the state evolution of (3.3) after t_2 . In other words, state information eventually propagates to all other users in the network and old information is eventually purged from the network. In the power control problem, Assumption 3.1 is always satisfied unless the communication between users is totally lost during a semi-infinite time interval.

Note that all bounded delays, irrespective of whether they are constant or time-varying, satisfy Assumption 3.1. Moreover, delays satisfying (3.2) may be unbounded. Consider the following particular class of unbounded delays which was studied in [26].

Assumption 3.2. There exist $T > 0$ and a scalar $0 \leq \alpha < 1$ such that

$$(3.4) \quad \sup_{t > T} \frac{\tau(t)}{t} = \alpha.$$

One can easily verify that (3.4) implies (3.2). However, the next example shows that the converse does not, in general, hold. Hence, Assumption 3.2 is a special case of Assumption 3.1.

Example 3.1. Let $\tau(t) = t - \ln(t + 1)$ for $t \geq 0$. Since

$$\begin{aligned} \lim_{t \rightarrow +\infty} t - \tau(t) &= \lim_{t \rightarrow +\infty} \ln(t + 1) = +\infty, \\ \lim_{t \rightarrow +\infty} \frac{\tau(t)}{t} &= \lim_{t \rightarrow +\infty} \frac{t - \ln(t + 1)}{t} = 1, \end{aligned}$$

it is clear that (3.2) holds while (3.4) does not hold.

Remark 3.1. Assumption 3.1 implies that there exists a sufficiently large $T_0 > 0$ such that $t - \tau(t) > 0$ for all $t > T_0$. Define

$$\tau_{\max} = - \inf_{0 \leq t \leq T_0} \left\{ t - \tau(t) \right\}.$$

Since $\tau_{\max} \geq 0$ is bounded, it follows that for any delay satisfying Assumption 3.1, even if it is unbounded, the initial condition $\varphi(\cdot)$ is defined on a bounded set $[-\tau_{\max}, 0]$.

In this section, we study delay-independent stability of nonlinear systems of the form (3.1) which are *positive* defined as follows.

DEFINITION 3.1. *System \mathcal{G} given by (3.1) is said to be positive if for every nonnegative initial condition $\varphi(\cdot) \in \mathcal{C}([-\tau_{\max}, 0], \mathbb{R}_+^n)$, the corresponding state trajectory is nonnegative, that is, $\mathbf{x}(t) \in \mathbb{R}_+^n$ for all $t \geq 0$.*

The following result provides a sufficient condition for positivity of \mathcal{G} .

PROPOSITION 3.2. *Consider the time-delay system \mathcal{G} given by (3.1). If the following condition holds,*

$$(3.5) \quad \begin{aligned} \forall i \in \{1, \dots, n\}, \forall \mathbf{x} \in \mathbb{R}_+^n : x_i = 0 \Rightarrow f_i(\mathbf{x}) \geq 0, \\ \forall \mathbf{x} \in \mathbb{R}_+^n, \quad \mathbf{g}(\mathbf{x}) \geq 0, \end{aligned}$$

then \mathcal{G} is positive.

Proof. See Appendix A. \square

Note that the nonnegativity of the initial condition is essential for ensuring positivity of the state evolution of the system \mathcal{G} given by (3.1). In other words, when $\varphi(\cdot) \geq \mathbf{0}$ is not satisfied, $\mathbf{x}(t)$ may not stay in the positive orthant even if the condition of Proposition 3.2 hold.

In [18, Proposition 3.1], it was shown that for nonzero constant delays, the sufficient condition in Proposition 3.2 is also necessary, i.e., a system \mathcal{G} given by (3.1) with $\tau(t) = \tau_{\max} > 0, t \geq 0$, is positive if and only if (3.5) holds. However, the condition is not necessary when we allow for time-varying delays, as the next example shows.

Example 3.2. Consider a continuous-time linear system described by (3.1) with

$$(3.6) \quad \mathbf{f}(x_1, x_2) = \begin{bmatrix} 1 & 0 \\ -1 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}, \quad \mathbf{g}(x_1, x_2) = \begin{bmatrix} 0 & 0 \\ e & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix},$$

where e is the base of the natural logarithm, and let the time-varying delay be

$$(3.7) \quad \tau(t) = \begin{cases} 0, & 0 \leq t \leq 1, \\ t - 1, & 1 \leq t \leq 2, \\ 1, & 2 \leq t. \end{cases}$$

Note that $0 \leq \tau(t) \leq 1$ for all $t \geq 0$. The solution to (3.6) is given by

$$\begin{aligned} x_1(t) &= x_1(0)e^t, & 0 \leq t, \\ x_2(t) &= \begin{cases} x_2(0) + (e - 1)(e^t - 1)x_1(0), & 0 \leq t \leq 1, \\ x_2(0) + (e^2t - e^t + 1 - e)x_1(0), & 1 \leq t \leq 2, \\ x_2(0) + (e^2 - e + 1)x_1(0), & 2 \leq t. \end{cases} \end{aligned}$$

It is straightforward to verify that $\mathbf{x}(t) \geq \mathbf{0}$ for every nonnegative initial condition $\mathbf{x}(0) = (x_1(0), x_2(0))$, and hence the linear system (3.6) with the bounded time-varying delay (3.7) is positive. However, the sufficient condition given in Proposition 3.2 is not satisfied in this example, since $x_2 = 0$ does not imply $f_2(\mathbf{x}) \geq 0$ for all $\mathbf{x} \in \mathbb{R}_+^2$ (take, for example, $f_2(1, 0) = -1 < 0$).

From this point on, vector fields \mathbf{f} and \mathbf{g} satisfy Assumption 3.3.

Assumption 3.3. The following properties hold:

1. \mathbf{f} is cooperative and \mathbf{g} is nondecreasing on \mathbb{R}_+^n .
2. \mathbf{f} and \mathbf{g} are homogeneous of degree p with respect to the dilation map $\delta_\lambda^r(\mathbf{x})$.

A system \mathcal{G} given by (3.1) satisfying Assumption 3.3 is called *homogeneous cooperative*. Since $\mathbf{f}(\mathbf{0}) = \mathbf{g}(\mathbf{0}) = \mathbf{0}$, by Propositions 2.2 and 3.2, Assumption 3.3.1 ensures the positivity of homogeneous cooperative systems. The model of some physical systems fall within this class of positive systems. For example, continuous-time linear and several nonlinear power control algorithms for wireless networks are described by homogeneous cooperative systems [3].

While the stability of general dynamical systems may depend on the magnitude and variation of the time delays, homogeneous cooperative systems have been shown to be insensitive to constant delays [4]. More precisely, the homogeneous cooperative system (3.1) with a constant delay $\tau(t) = \tau_{\max}$, $t \geq 0$, is globally asymptotically stable for all $\tau_{\max} \geq 0$ if and only if the undelayed system ($\tau_{\max} = 0$) is globally asymptotically stable. The main goal of this section is to (i) determine whether a similar delay-independent stability result holds for homogeneous cooperative systems with time-varying delays satisfying Assumption 3.1 and to (ii) give explicit estimates of the decay rate for different classes of time delays (e.g., bounded delays, unbounded delays satisfying Assumption 3.2, etc.).

3.2. Asymptotic stability of homogeneous cooperative systems. The following theorem establishes a necessary and sufficient condition for delay-independent asymptotic stability of homogeneous cooperative systems with time-varying delays satisfying Assumption 3.1. Our proof (which is conceptually related to the Lyapunov stability theorem) uses the Lyapunov function

$$V(\mathbf{x}) = \max_{1 \leq i \leq n} \left(\frac{x_i}{v_i} \right)^{\frac{r_{\max}}{r_i}},$$

where $\mathbf{v} > \mathbf{0}$, and $r_{\max} = \max_{1 \leq i \leq n} r_i$, to define sets

$$(3.8) \quad S(m) = \left\{ \mathbf{x} \in \mathbb{R}_+^n \mid V(\mathbf{x}) \leq \gamma^m \|\boldsymbol{\varphi}\| \right\}, \quad m \in \mathbb{N}_0,$$

where $0 \leq \gamma < 1$, and

$$(3.9) \quad \|\boldsymbol{\varphi}\| = \sup_{-\tau_{\max} \leq s \leq 0} V(\boldsymbol{\varphi}(s)),$$

and shows that for each m , there exists $t_m \geq 0$ such that $\mathbf{x}(t) \in S(m)$ for all $t \geq t_m$. In other words, the system state will enter each set $S(m)$ at some time t_m and remain in the set for all future times. Since the sets are nested, i.e.,

$$S(0) \supset \cdots \supset S(m) \supset S(m+1) \supset \cdots,$$

the state will move sequentially from set $S(m)$ to $S(m+1)$; cf. Figure 1. Thus, the sets play a similar role as level sets of the Lyapunov function $V(\mathbf{x})$. Note that when \mathbf{f} and \mathbf{g} are homogeneous with respect to the standard dilation map, $V(\mathbf{x}) = \|\mathbf{x}\|_\infty^p$, which is often used in analysis of positive linear systems [38].

THEOREM 3.3. *For the homogeneous cooperative system \mathcal{G} given by (3.1), suppose that Assumption 3.1 holds. Then, the following statements are equivalent:*




FIG. 1. Level curves of the Lyapunov function $V(\mathbf{x})$ in the two-dimensional case. The key idea behind the proof of Theorem 3.3 is that $\varphi(\cdot)$ is initially within the set $S(0)$ and at some time $t_1 \geq 0$ eventually $\mathbf{x}(t)$ enters and stays within the set $S(1)$ for all $t \geq t_1$; moreover, as t increases further, $\mathbf{x}(t)$ sequentially moves into other sets.

(i) There exists a vector $\mathbf{v} > \mathbf{0}$ such that

$$(3.10) \quad \mathbf{f}(\mathbf{v}) + \mathbf{g}(\mathbf{v}) < \mathbf{0}.$$

(ii) The positive system \mathcal{G} is globally asymptotically stable for every nonnegative initial condition $\varphi(\cdot) \in \mathcal{C}([-\tau_{\max}, 0], \mathbb{R}_+^n)$ and for all time delays satisfying Assumption 3.1.

(iii) The positive system \mathcal{G} without delay ($\tau(t) = 0, t \geq 0$) is globally asymptotically stable for all nonnegative initial conditions.

Proof. See Appendix B. \square

The stability condition (3.10) does not include any information on the magnitude and variation of delays, so it ensures delay-independent stability. According to Theorem 3.3, the homogeneous cooperative system \mathcal{G} given by (3.1) is globally asymptotically stable for all time delays satisfying Assumption 3.1 if and only if the corresponding delay-free system is globally asymptotically stable. This property is very useful in practical applications, since the delays may not be easy to model in detail.

Note that Theorem 3.3 can be easily extended to homogeneous cooperative systems with multiple delays of the form

$$\dot{\mathbf{x}}(t) = \mathbf{f}(\mathbf{x}(t)) + \sum_{q=1}^s \mathbf{g}_q(\mathbf{x}(t - \tau_q(t))).$$

Here, $s \in \mathbb{N}$, \mathbf{f} is cooperative and homogeneous, \mathbf{g}_q for $q = 1, \dots, s$ are homogeneous and nondecreasing on \mathbb{R}_+^n , and $\tau_q(t)$ satisfy Assumption 3.1. In this case, the stability condition (3.10) becomes

$$\mathbf{f}(\mathbf{v}) + \sum_{q=1}^s \mathbf{g}_q(\mathbf{v}) < \mathbf{0}.$$

Remark 3.2. In [13], it has been shown that if \mathbf{f} and \mathbf{g} are homogeneous of degree zero with respect to the standard dilation map, the homogeneous cooperative system (3.1) is insensitive to bounded time-varying delays. In this work, we extend

this result to cooperative systems that are homogeneous of any degree with respect to an arbitrary dilation map. Moreover, we impose minimal restrictions on time delays and establish insensitivity of homogeneous cooperative systems to the general class of delays described by Assumption 3.1, which includes bounded delays as a special case.

3.3. Decay rates of homogeneous cooperative systems. Theorem 3.3 is concerned with the asymptotic stability of homogeneous cooperative systems with time-varying delays. However, there are processes and applications for which it is desirable that the system has a certain decay rate. Loosely speaking, the system has to converge quickly enough to the equilibrium. Hence, it is important to investigate the impact of delays on the decay rate of such systems. In this section, we characterize how time delays affect the decay rate of the homogeneous cooperative system \mathcal{G} given by (3.1). Before stating the main result, we provide the definition of μ -stability, introduced in [7], for continuous-time systems.

DEFINITION 3.4 (see [7]). *Suppose that $\mu : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is a nondecreasing function satisfying $\mu(t) \rightarrow +\infty$ as $t \rightarrow +\infty$. System \mathcal{G} given by (3.1) is said to be globally μ -stable if there exists a constant $M > 0$ such that for any initial function $\varphi(\cdot)$, the solution $\mathbf{x}(t)$ satisfies*

$$\|\mathbf{x}(t)\| \leq \frac{M}{\mu(t)}, \quad t > 0,$$

where $\|\cdot\|$ is some norm on \mathbb{R}^n .

This definition can be regarded as a unification of several types of stability. For example, when $\mu(t) = e^{\eta t}$ with $\eta > 0$, the μ -stability becomes *exponential stability*; and when $\mu(t) = t^\xi$ with $\xi > 0$, then the μ -stability becomes *power-rate stability*.

Global μ -stability of homogenous cooperative systems can be verified using the following theorem.

THEOREM 3.5. *Consider the homogeneous cooperative system \mathcal{G} given by (3.1). Suppose that Assumption 3.1 holds, and that there is a vector $\mathbf{v} > \mathbf{0}$ satisfying*

$$(3.11) \quad \mathbf{f}(\mathbf{v}) + \mathbf{g}(\mathbf{v}) < \mathbf{0}.$$

If there exists a function $\mu : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ such that the following conditions hold,

- (i) $\mu(t) > 0$ for all $t > 0$,
- (ii) $\mu(t)$ is a nondecreasing function,
- (iii) $\lim_{t \rightarrow +\infty} \mu(t) = +\infty$,
- (iv) for each $i \in \{1, \dots, n\}$,

$$\left(\frac{r_{\max}}{r_i} \right) \left(\left(\frac{f_i(\mathbf{v})}{v_i} \right) + \left(\lim_{t \rightarrow \infty} \frac{\mu(t)}{\mu(t - \tau(t))} \right)^{\frac{r_i + p}{r_{\max}}} \left(\frac{g_i(\mathbf{v})}{v_i} \right) \right) + \lim_{t \rightarrow \infty} \frac{\dot{\mu}(t)}{(\mu(t))^{1 - \frac{p}{r_{\max}}}} < 0,$$

then every solution of \mathcal{G} starting in the positive orthant satisfies

$$\left(\frac{x_i(t)}{v_i} \right)^{\frac{r_{\max}}{r_i}} = O(\mu^{-1}(t)), \quad t \geq 0,$$

for each $i = 1, \dots, n$.

Proof. See Appendix C. \square

According to Theorem 3.5, any function $\mu(t)$ satisfying conditions (i)–(iv) can be used to estimate the decay rate of homogeneous cooperative systems with time-varying delays satisfying Assumption 3.1. From condition (iv), it is clear that the asymptotic behavior of the delay $\tau(t)$ influences the admissible choices for $\mu(t)$ and, hence, the decay bounds that we are able to guarantee. To clarify this statement, we will analyze a few special cases in detail. First, assume that $\tau(t)$ is bounded, i.e.,

$$(3.12) \quad 0 \leq \tau(t) \leq \tau_{\max}, \quad t \geq 0.$$

The following result shows that the decay rate of homogeneous cooperative systems of degree p with bounded time-varying delays is upper bounded by an exponential function of time when $p = 0$ and by a polynomial function of time when $p > 0$.

COROLLARY 3.6. *For the homogeneous cooperative system \mathcal{G} given by (3.1), suppose that (3.12) holds and that there exists a vector $\mathbf{v} > \mathbf{0}$ satisfying (3.11).*

(i) *If \mathbf{f} and \mathbf{g} are homogeneous of degree $p = 0$, then \mathcal{G} is globally exponentially stable. In particular,*

$$(3.13) \quad \left(\frac{x_i(t)}{v_i} \right)^{\frac{r_{\max}}{r_i}} = O(e^{-\eta t}), \quad t \geq 0,$$

where $0 < \eta < \min_{1 \leq i \leq n} \eta_i$, and η_i is the positive solution of the equation

$$(3.14) \quad \left(\frac{r_{\max}}{r_i} \right) \left(\left(\frac{f_i(\mathbf{v})}{v_i} \right) + \left(e^{\eta_i \tau_{\max}} \right)^{\frac{r_i}{r_{\max}}} \left(\frac{g_i(\mathbf{v})}{v_i} \right) \right) + \eta_i = 0;$$

(ii) *if \mathbf{f} and \mathbf{g} are homogeneous of degree $p > 0$, the solution $\mathbf{x}(t)$ of \mathcal{G} satisfies*

$$(3.15) \quad \left(\frac{x_i(t)}{v_i} \right)^{\frac{r_{\max}}{r_i}} = O\left((\theta t + 1)^{-\frac{r_{\max}}{p}} \right), \quad t \geq 0,$$

where $0 < \theta < \min\left\{ \frac{1}{\tau_{\max}}, \min_{1 \leq i \leq n} \theta_i \right\}$, and θ_i is the positive solution to

$$(3.16) \quad \frac{f_i(\mathbf{v})}{v_i} + \frac{g_i(\mathbf{v})}{v_i} + \theta_i \frac{r_i}{p} = 0.$$

Proof. See Appendix D. □

Remark 3.3. Equation (3.14) has three parameters: the maximum delay bound τ_{\max} , the positive vector \mathbf{v} , and η_i . For any fixed $\tau_{\max} \geq 0$ and any fixed $\mathbf{v} > \mathbf{0}$ satisfying (3.11), the left-hand side of (3.14) is smaller than the right-hand side for $\eta_i = 0$ and strictly monotonically increasing in $\eta_i > 0$. Therefore, (3.14) has always a unique positive solution η_i . By a similar argument, (3.16) also admits a unique positive solution θ_i .

While the stability of homogeneous cooperative systems with delays satisfying Assumption 3.1 may, in general, only be asymptotic, Corollary 3.6 demonstrates that if the delays are bounded, we can guarantee certain decay rates. We will now establish similar decay bounds for unbounded delays satisfying Assumption 3.2.

COROLLARY 3.7. *Consider the homogeneous cooperative system \mathcal{G} given by (3.1). Suppose that Assumption 3.2 holds and that there is a vector $\mathbf{v} > \mathbf{0}$ satisfying (3.11). Then, \mathcal{G} is globally power-rate stable. In particular,*

(i) *if \mathbf{f} and \mathbf{g} are homogeneous of degree $p = 0$, the solution $\mathbf{x}(t)$ of \mathcal{G} satisfies*

$$\left(\frac{x_i(t)}{v_i} \right)^{\frac{r_{\max}}{r_i}} = O(t^{-\xi}), \quad t \geq 0,$$




FIG. 2. Plot of β for different values of the parameter $\alpha \in [0, 1)$. Clearly, β is monotonically decreasing with α and approaches zero as α tends to one.

where $0 < \xi < \min_{1 \leq i \leq n} \xi_i$, and ξ_i is the unique positive solution to

$$(3.17) \quad \left(\frac{f_i(\mathbf{v})}{v_i} \right) + \left(\frac{1}{1-\alpha} \right)^{\frac{r_i}{r_{\max}} \xi_i} \left(\frac{g_i(\mathbf{v})}{v_i} \right) = 0;$$

(ii) if \mathbf{f} and \mathbf{g} are homogeneous of degree $p > 0$, then

$$\left(\frac{x_i(t)}{v_i} \right)^{\frac{r_{\max}}{r_i}} = O\left(t^{-\frac{r_{\max}}{p} \beta}\right), \quad t \geq 0,$$

where $\beta \in (0, 1)$ is such that

$$(3.18) \quad \left(\frac{f_i(\mathbf{v})}{v_i} \right) + \left(\frac{1}{1-\alpha} \right)^{\left(1 + \frac{r_i}{p}\right)\beta} \left(\frac{g_i(\mathbf{v})}{v_i} \right) < 0$$

holds for all i .

Proof. See Appendix E. \square

Corollary 3.7 shows that the decay rate of homogeneous cooperative systems of degree zero with unbounded delays satisfying Assumption 3.2 is of order $O(t^{-\xi})$. Equation (3.17) quantifies how the magnitude of the upper delay bound, α , affects ξ . Specifically, ξ_i is monotonically decreasing with α and approaches zero as α tends to one. By similar reasoning, β , on which the guaranteed decay rate of homogeneous cooperative systems of degree greater than zero depends, in (3.18) approaches zero as α tends to one (see Figure 2). Hence, while the homogeneous cooperative system (3.1) remains power-rate stable for arbitrary unbounded delays satisfying Assumption 3.2, the decay rate deteriorates with increasing α . This means that the guaranteed convergence rates get increasingly slower as delays are allowed to grow quicker when $t \rightarrow \infty$.

3.4. A special case: Continuous-time positive linear systems. Let $\mathbf{f}(\mathbf{x}) = A\mathbf{x}$ and $\mathbf{g}(\mathbf{x}) = B\mathbf{x}$ such that $A \in \mathbb{R}^{n \times n}$ is Metzler and $B \in \mathbb{R}^{n \times n}$ is nonnegative. Then, the homogeneous cooperative system (3.1) reduces to the positive linear system

$$(3.19) \quad \mathcal{G}_L : \begin{cases} \dot{\mathbf{x}}(t) = A\mathbf{x}(t) + B\mathbf{x}(t - \tau(t)), & t \geq 0, \\ \mathbf{x}(t) = \boldsymbol{\varphi}(t), & t \in [-\tau_{\max}, 0]. \end{cases}$$

We then have the following special case of Theorem 3.3.

COROLLARY 3.8. Consider the positive linear system \mathcal{G}_L given by (3.19), where A is Metzler and B is nonnegative. Then, \mathcal{G}_L is globally asymptotically stable for all time delays satisfying Assumption 3.1 if and only if there exists a vector $\mathbf{v} > \mathbf{0}$ such that

$$(3.20) \quad (A + B)\mathbf{v} < \mathbf{0}.$$

Corollary 3.8 shows that if the positive linear system (3.19) without delay is stable, it remains asymptotically stable under all bounded and unbounded time-varying delays satisfying Assumption 3.1. Note that the stability condition (3.20) is a linear programming feasibility problem in \mathbf{v} which can be verified numerically in polynomial time.

Remark 3.4. Since A is Metzler and B is nonnegative, $A + B$ is Metzler. It follows from [38, Proposition 2] that the linear inequality (3.20) holds if and only if $A + B$ is Hurwitz, i.e., all its eigenvalues have negative real parts.

While the asymptotic stability of the positive linear system \mathcal{G}_L given by (3.19) with time-varying delays satisfying Assumption 3.1 has been investigated in [42], the impact of time delays on the decay rate has been missing. Theorem 3.5 helps us to find guaranteed decay rates of \mathcal{G}_L for different classes of time delays. Specifically, Corollaries 3.6 and 3.7 show that \mathcal{G}_L is exponentially stable if time-varying delays are bounded, and power-rate stable if delays are unbounded and satisfy Assumption 3.2. Therefore, not only do we extend the result of [42] to general homogeneous cooperative systems (not necessarily linear), but we also provide explicit bounds on the decay rate of positive linear systems.

Remark 3.5. In [27, Example 4.5], it was shown that a positive linear system with unbounded delays satisfying Assumption 3.2 may converge slower than any exponential function. However, an upper bound for the decay rate was not derived in [27]. Corollary 3.7 reveals that under Assumption 3.2 on delays, the decay rate of positive linear systems is upper bounded by a polynomial function of time.

4. Discrete-time homogeneous nondecreasing systems.

4.1. Problem statement. Next, we consider the discrete-time analogue of (3.1):

$$(4.1) \quad \Sigma : \begin{cases} \mathbf{x}(k + 1) = \mathbf{f}(\mathbf{x}(k)) + \mathbf{g}(\mathbf{x}(k - d(k))), & k \in \mathbb{N}_0, \\ \mathbf{x}(k) = \boldsymbol{\varphi}(k), & k \in \{-d_{\max}, \dots, 0\}. \end{cases}$$

Here, $\mathbf{x}(k) \in \mathbb{R}^n$ is the state variable, $\mathbf{f}, \mathbf{g} : \mathbb{R}^n \rightarrow \mathbb{R}^n$ are continuous vector fields with $\mathbf{f}(\mathbf{0}) = \mathbf{g}(\mathbf{0}) = \mathbf{0}$, $d_{\max} \in \mathbb{N}_0$, $\boldsymbol{\varphi} : \{-d_{\max}, \dots, 0\} \rightarrow \mathbb{R}^n$ is the vector sequence specifying the initial state of the system, and $d(k)$ represents the time-varying delay which satisfies the following assumption.

Assumption 4.1. The delay $d : \mathbb{N}_0 \rightarrow \mathbb{N}_0$ satisfies

$$(4.2) \quad \lim_{k \rightarrow +\infty} k - d(k) = +\infty.$$

Intuitively, if Assumption 4.1 does not hold, computation of $\mathbf{x}(k)$, even for large values of k , may involve the initial condition $\boldsymbol{\varphi}(\cdot)$ and those states near it, and hence $\mathbf{x}(k)$ may not converge to zero as $k \rightarrow \infty$. To avoid this situation, Assumption 4.1 guarantees that old state information is eventually not used in evaluating (4.1).

Remark 4.1. Assumption 4.1 implies that there exists a sufficiently large $T_0 \in \mathbb{N}_0$ such that $k - d(k) > 0$ for all $k > T_0$. Let

$$d_{\max} = - \inf_{0 \leq k \leq T_0} \left\{ k - d(k) \right\}.$$

Clearly, $d_{\max} \in \mathbb{N}_0$ is bounded. It follows that, even for unbounded delays satisfying Assumption 4.1, the initial condition $\varphi(\cdot)$ is defined on a finite set $\{-d_{\max}, \dots, 0\}$.

DEFINITION 4.1. *The system Σ given by (4.1) is said to be positive if for every nonnegative initial condition $\varphi(\cdot) \in \mathbb{R}_+^n$, the corresponding solution is nonnegative, that is, $\mathbf{x}(k) \geq \mathbf{0}$ for all $k \in \mathbb{N}$.*

Positivity of Σ is readily verified using the following result.

PROPOSITION 4.2. *Consider the time-delay system Σ given by (4.1). If $\mathbf{f}(\mathbf{x}) \geq \mathbf{0}$ and $\mathbf{g}(\mathbf{x}) \geq \mathbf{0}$ for all $\mathbf{x} \in \mathbb{R}_+^n$, then Σ is positive.*

For nonzero constant delays ($d(k) = d_{\max} > 0$, $k \in \mathbb{N}_0$), the sufficient condition in Proposition 4.2 is also necessary [18, Proposition 3.4]. However, the following example shows that this result may not be true when delays are time-varying.

Example 4.1. Consider a discrete-time linear system described by (4.1) with

$$f(x) = 2x, \quad g(x) = -x, \quad d(k) = \frac{1}{2}(1 - (-1)^k), \quad k \in \mathbb{N}_0.$$

Since $g(x) < 0$ for $x > 0$, the sufficient condition given in Proposition 4.2 is not satisfied. However, it is easy to verify that the solution of this system is $x(k) = x(0)$, $k \in \mathbb{N}_0$, and hence $x(k) \geq 0$ for all $x(0) \geq 0$.

In this section, vector fields \mathbf{f} and \mathbf{g} satisfy the next assumption.

Assumption 4.2. The following properties hold:

1. \mathbf{f} and \mathbf{g} are nondecreasing on \mathbb{R}_+^n .
2. \mathbf{f} and \mathbf{g} are homogeneous of degree p with respect to the dilation map $\delta_\lambda^r(\mathbf{x})$.

A system Σ given by (4.1) satisfying Assumption 4.2 is called *homogeneous nondecreasing*. Since $\mathbf{f}(\mathbf{0}) = \mathbf{g}(\mathbf{0}) = \mathbf{0}$, Assumption 4.2.1 implies that \mathbf{f} and \mathbf{g} satisfy the condition of Proposition 4.2. Hence, homogeneous nondecreasing systems are positive.

Our main objective in this section is to study delay-independent stability of homogeneous nondecreasing systems of the form (4.1) with time-varying delays satisfying Assumption 4.1.

4.2. Asymptotic stability of homogeneous nondecreasing systems. The next theorem shows that global asymptotic stability of nondecreasing systems of the form (4.1) that are homogeneous of degree zero is insensitive to bounded and unbounded time-varying delays satisfying Assumption 4.1.

THEOREM 4.3. *For the homogeneous nondecreasing system Σ given by (4.1), suppose that \mathbf{f} and \mathbf{g} are homogeneous of degree $p = 0$. Then, the following statements are equivalent:*

- (i) *There exists a vector $\mathbf{v} > \mathbf{0}$ such that*

$$(4.3) \quad \mathbf{f}(\mathbf{v}) + \mathbf{g}(\mathbf{v}) < \mathbf{v}.$$

- (ii) Σ is globally asymptotically stable for any nonnegative initial conditions and for all bounded and unbounded time-varying delays satisfying Assumption 4.1.
- (iii) Σ without delay ($d(k) = 0$, $k \in \mathbb{N}_0$) is globally asymptotically stable for any nonnegative initial conditions.

Proof. See Appendix F. \square

Theorem 4.3 provides a test for global asymptotic stability of homogeneous nondecreasing systems of degree zero; if we can demonstrate the existence of a vector $\mathbf{v} > \mathbf{0}$ satisfying (4.3), then the origin is globally asymptotically stable for all delays satisfying Assumption 4.1. However, the following example illustrates that (4.3) is,

in general, not a sufficient condition for global asymptotic stability of homogeneous nondecreasing systems of degree greater than zero.

Example 4.2. Consider a discrete-time system described by (4.1) with $f(x) = x^2$ and $g(x) = 0$. Clearly, f is nondecreasing on \mathbb{R}_+ and homogeneous of degree one with respect to the standard dilation map. Since $f(0.5) = 0.25 < 0.5$, (4.3) holds. However, it is easy to verify that solutions of this system starting from initial conditions $x(0) \geq 1$ do not converge to the origin, i.e., the origin is not globally asymptotically stable.

We now show that under stability condition (4.3), homogeneous nondecreasing systems of degree greater than zero with time-varying delays satisfying Assumption 4.1 have a locally asymptotically stable equilibrium point at the origin, i.e., $\mathbf{x}(k)$ converges to the origin as $k \rightarrow \infty$ for sufficiently small initial conditions.

COROLLARY 4.4. *For the homogeneous nondecreasing system Σ given by (4.1) with degree $p > 0$, suppose that Assumption 4.1 holds. If there exists a vector $\mathbf{v} > \mathbf{0}$ such that (4.3) holds, then the origin is asymptotically stable with respect to initial conditions satisfying*

$$\mathbf{0} \leq \boldsymbol{\varphi}(k) \leq \mathbf{v} \quad \forall k \in \{-d_{\max}, \dots, 0\}.$$

Proof. See Appendix G. \square

4.3. Decay rates of homogeneous nondecreasing systems of degree zero.

The next definition introduces μ -stability for discrete-time systems.

DEFINITION 4.5. *Suppose that $\mu : \mathbb{N} \rightarrow \mathbb{R}_+$ is a nondecreasing function satisfying $\mu(k) \rightarrow +\infty$ as $k \rightarrow +\infty$. The system Σ given by (4.1) is said to be globally μ -stable if there exists a constant $M > 0$ such that for any initial function $\boldsymbol{\varphi}(\cdot)$, the solution $\mathbf{x}(k)$ satisfies*

$$\|\mathbf{x}(k)\| \leq \frac{M}{\mu(k)}, \quad k \in \mathbb{N},$$

where $\|\cdot\|$ is some norm on \mathbb{R}^n .

Paralleling our continuous-time results, global μ -stability of homogeneous nondecreasing systems of degree zero with time-varying delays can be established using the following theorem.

THEOREM 4.6. *Consider the homogeneous nondecreasing system Σ given by (4.1) with degree $p = 0$. Suppose that Assumption 4.1 holds, and that there is a vector $\mathbf{v} > \mathbf{0}$ satisfying*

$$(4.4) \quad \mathbf{f}(\mathbf{v}) + \mathbf{g}(\mathbf{v}) < \mathbf{v}.$$

If there exists a function $\mu : \mathbb{N} \rightarrow \mathbb{R}_+$ such that the following conditions hold,

- (i) $\mu(k) > 0$ for all $k \in \mathbb{N}$,
- (ii) $\mu(k + 1) \geq \mu(k)$ for all $k \in \mathbb{N}$,
- (iii) $\lim_{k \rightarrow +\infty} \mu(k) = +\infty$,
- (iv) for each $i \in \{1, \dots, n\}$,

$$\left(\lim_{k \rightarrow \infty} \frac{\mu(k + 1)}{\mu(k)} \right)^{\frac{r_i}{r_{\max}}} \left(\frac{f_i(\mathbf{v})}{v_i} \right) + \left(\lim_{k \rightarrow \infty} \frac{\mu(k + 1)}{\mu(k - d(k))} \right)^{\frac{r_i}{r_{\max}}} \left(\frac{g_i(\mathbf{v})}{v_i} \right) < 1,$$

then every solution of Σ starting in the positive orthant satisfies

$$\left(\frac{x_i(k)}{v_i} \right)^{\frac{r_{\max}}{r_i}} = O(\mu^{-1}(k)), \quad k \in \mathbb{N},$$

for each $i = 1, \dots, n$.

Proof. See Appendix H. \square

Theorem 4.6 allows us to establish convergence rates of homogeneous nondecreasing systems of degree zero under various classes of time-varying delays. We give the following result.

COROLLARY 4.7. *For the homogeneous nondecreasing system Σ given by (4.1) with degree $p = 0$, suppose that there exists a vector $\mathbf{v} > \mathbf{0}$ satisfying (4.4), and that there exist $T \in \mathbb{N}$ and a scalar $0 \leq \alpha < 1$ such that*

$$(4.5) \quad \sup_{k>T} \frac{d(k)}{k} = \alpha.$$

Let ξ_i be the unique positive solution of the equation

$$(4.6) \quad \left(\frac{f_i(\mathbf{v})}{v_i} \right) + \left(\frac{1}{1-\alpha} \right)^{\frac{r_i}{r_{\max}} \xi_i} \left(\frac{g_i(\mathbf{v})}{v_i} \right) = 1, \quad i = 1, \dots, n.$$

Then, Σ is globally power-rate stable for any nonnegative initial conditions and for any unbounded delays satisfying (4.5). In particular,

$$\left(\frac{x_i(k)}{v_i} \right)^{\frac{r_{\max}}{r_i}} = O(k^{-\xi}), \quad k \in \mathbb{N},$$

where $0 < \xi < \min_{1 \leq i \leq n} \xi_i$.

4.4. A special case: Discrete-time positive linear systems. We now discuss delay-independent stability of a special case of (4.1), namely, discrete-time positive linear systems of the form

$$(4.7) \quad \Sigma_L : \begin{cases} \mathbf{x}(k+1) = A\mathbf{x}(k) + B\mathbf{x}(k-d(k)), & k \in \mathbb{N}_0, \\ \mathbf{x}(k) = \boldsymbol{\varphi}(k), & k \in \{-d_{\max}, \dots, 0\}. \end{cases}$$

In terms of (4.1), $\mathbf{f}(\mathbf{x}) = A\mathbf{x}$ and $\mathbf{g}(\mathbf{x}) = B\mathbf{x}$. It is easy to verify that if $A, B \in \mathbb{R}^{n \times n}$ are nonnegative matrices, Assumption 4.2 is satisfied. Therefore, Theorem 4.3 can help us to derive a necessary and sufficient condition for delay-independent stability of (4.7). Specifically, we note the following.

COROLLARY 4.8. *Consider the discrete-time positive linear system Σ_L given by (4.7), where A and B are nonnegative. Then, there exists a vector $\mathbf{v} > \mathbf{0}$ such that*

$$(4.8) \quad (A+B)\mathbf{v} < \mathbf{v}$$

if and only if Σ_L is globally asymptotically stable for all time delays satisfying Assumption 4.1.

Remark 4.2. For the positive linear system (4.7), A and B are nonnegative, so $A+B$ is also nonnegative. According to property of nonnegative matrices [2], [38, Proposition 1], there exists a vector $\mathbf{v} > \mathbf{0}$ satisfying (4.8) if and only if all eigenvalues of $A+B$ are strictly inside the unit circle.

Remark 4.3. In [12], it was shown that discrete-time positive linear systems are insensitive to time delays satisfying Assumption 4.1. Theorem 4.3 shows that a similar delay-independent stability result holds for nonlinear positive systems whose vector fields are nondecreasing and homogeneous of degree zero. Furthermore, the impact of various classes of time delays on the convergence rate of positive linear systems has been missing in [12], whereas Theorem 4.6 provides explicit bounds on the decay rate that allow us to quantify the impact of bounded and unbounded time-varying delays on the decay rate.




FIG. 3. Comparison of guaranteed upper bound and actual decay rate of the homogeneous cooperative system (5.1) corresponding to the initial condition $\varphi(t) = (1, 1)$ for all $t \in [-5, 0]$.

5. An illustrative example. Consider the continuous-time system (3.1) with

$$(5.1) \quad \mathbf{f}(x_1, x_2) = \begin{bmatrix} -5x_1^3 + 2x_1x_2 \\ x_1^2x_2 - 4x_2^2 \end{bmatrix}, \quad \mathbf{g}(x_1, x_2) = \begin{bmatrix} x_1x_2 \\ 2x_1^4 \end{bmatrix}.$$

Both \mathbf{f} and \mathbf{g} are homogeneous of degree $p = 2$ with respect to the dilation map $\delta_\lambda^r(\mathbf{x})$ with $\mathbf{r} = (1, 2)$. Moreover, \mathbf{f} is cooperative and \mathbf{g} is nondecreasing on \mathbb{R}_+^2 . Since $\mathbf{f}(1, 1) + \mathbf{g}(1, 1) < \mathbf{0}$, it follows from Theorem 3.3 that the homogeneous cooperative system (5.1) is globally asymptotically stable for any time delays satisfying Assumption 3.1. Now, consider the specific time-varying delay $\tau(t) = 4 + \sin(t)$, $t \geq 0$. As $\tau(t) \leq \tau_{\max} = 5$ for all $t \geq 0$, Corollary 3.6 can help us to calculate an upper bound on the decay rate of (5.1). Using $\mathbf{v} = (1, 1)$ and $r_{\max} = 2$, the solutions to (3.16) are $\theta_1 = 4$, $\theta_2 = 1$, which implies that

$$\theta \cong \min \left\{ \frac{1}{5}, \min\{4, 1\} \right\} = \frac{1}{5}.$$

Thus, the solution $\mathbf{x}(t)$ of (5.1) satisfies

$$\max\{x_1^2(t), x_2(t)\} = O\left(\frac{1}{\frac{1}{5}t + 1}\right), \quad t \geq 0.$$

Figure 3 gives the simulation results of the actual decay rate of the homogeneous cooperative system (5.1) and the guaranteed decay rate we calculated, when the initial condition is $\varphi(t) = (1, 1)$ for all $t \in [-5, 0]$.

6. Conclusions. This paper has been concerned with delay-independent stability of a significant class of nonlinear (continuous- and discrete-time) positive systems with time-varying delays. We derived a set of necessary and sufficient conditions for global asymptotic stability of continuous-time homogeneous cooperative systems of arbitrary degree and discrete-time homogeneous nondecreasing systems of degree zero with bounded and unbounded time-varying delays. These results show that the asymptotic stability of such systems is independent of the magnitude and variation of the time delays. However, we also observed that the decay rates of these systems depend on how fast the delays can grow large. We developed two theorems for global μ -stability of positive systems that quantify the convergence rates for various classes of time delays. For discrete-time homogeneous nondecreasing systems of degree greater than zero, we demonstrated that the origin is locally asymptotically stable under global asymptotic stability conditions that we derived.

Appendix A. Proof of Proposition 3.2. Consider the following differential equation:

$$(A.1) \quad \begin{cases} \dot{\mathbf{y}}(t) = \mathbf{f}(\mathbf{y}(t)) + \mathbf{g}(\mathbf{y}(t - \tau(t))) + \frac{1}{k}\mathbf{1}, & t \geq 0, \\ \mathbf{y}(t) = \boldsymbol{\varphi}(t), & t \in [-\tau_{\max}, 0], \end{cases}$$

where $k \in \mathbb{N}$, and $\mathbf{1} \in \mathbb{R}^n$ is the vector with all components equal to 1. Let $\mathbf{y}^{(k)}(t)$ be the solution to (A.1) with the nonnegative initial condition $\boldsymbol{\varphi}(\cdot) \in \mathcal{C}([-\tau_{\max}, 0], \mathbb{R}_+^n)$. Clearly, $\mathbf{y}^{(k)}(0) = \boldsymbol{\varphi}(0) \geq \mathbf{0}$. We claim that $\mathbf{y}^{(k)}(t) \geq \mathbf{0}$ for all $t \geq 0$. By contradiction, suppose this is not true. Then, there exist an index $j \in \{1, \dots, n\}$ and a time $t_1 \geq 0$ such that $\mathbf{y}^{(k)}(t) \geq \mathbf{0}$ for all $t \in [0, t_1]$, $y_j^{(k)}(t_1) = 0$, and

$$(A.2) \quad D^+ y_j^{(k)}(t) \Big|_{t=t_1} \leq 0.$$

It follows from (3.5) and the above observations that

$$(A.3) \quad f_j(\mathbf{y}^{(k)}(t_1)) \geq 0.$$

Since $t_1 - \tau(t_1) \in [-\tau_{\max}, t_1]$ and $\boldsymbol{\varphi}(\cdot) \geq \mathbf{0}$, we have $\mathbf{y}^{(k)}(t_1 - \tau(t_1)) \geq \mathbf{0}$ irrespective of whether $t_1 - \tau(t_1)$ is nonnegative or not. From (3.5), we then have

$$(A.4) \quad g_j(\mathbf{y}^{(k)}(t_1 - \tau(t_1))) \geq 0.$$

Using (A.3) and (A.4), the upper-right Dini-derivative of $y_j^{(k)}(t)$ along the trajectories of (A.1) at $t = t_1$ is given by

$$\begin{aligned} D^+ y_j^{(k)}(t) \Big|_{t=t_1} &= f_j(\mathbf{y}^{(k)}(t_1)) + g_j(\mathbf{y}^{(k)}(t_1 - \tau(t_1))) + \frac{1}{k} \\ &\geq \frac{1}{k} \\ &> 0, \end{aligned}$$

which contradicts (A.2). Therefore,

$$(A.5) \quad \mathbf{y}^{(k)}(t) \geq \mathbf{0} \quad \forall t \geq 0.$$

As k was an arbitrary natural number, it follows that (A.5) holds for all $k \in \mathbb{N}$. By letting $k \rightarrow \infty$, $\mathbf{y}^{(k)}(t)$ converges to the solution $\mathbf{x}(t)$ of (3.1) uniformly on $[-\tau_{\max}, \infty)$

[21, Theorem 2.1], which implies that $\mathbf{x}(t) \geq \mathbf{0}$ for all $t \geq 0$. Hence, the system (3.1) is positive.

Appendix B. Proof of Theorem 3.3. (i) \Rightarrow (ii). Let $\mathbf{v} > \mathbf{0}$ be a vector such that (3.10) holds and let

$$(B.1) \quad \zeta = - \max_{1 \leq i \leq n} \left\{ \frac{f_i(\mathbf{v}) + g_i(\mathbf{v})}{v_i} \right\}.$$

Note that $\zeta > 0$. Since \mathbf{g} is nondecreasing on \mathbb{R}_+^n and $\mathbf{g}(\mathbf{0}) = \mathbf{0}$, $g_i(\mathbf{v}) \geq 0$ for all i . Thus,

$$(B.2) \quad \frac{f_i(\mathbf{v})}{v_i} \leq -\zeta$$

for $i = 1, \dots, n$. Define

$$\gamma_i = \left(1 + \frac{\zeta/2}{f_i(\mathbf{v})/v_i} \right)^{\frac{r_{\max}}{r_i+p}},$$

where $r_{\max} = \max_{1 \leq i \leq n} r_i$. From (B.2), one can verify that $\gamma_i \in (0, 1)$. We have

$$\begin{aligned} \gamma_i^{\frac{r_i+p}{r_{\max}}} \left(\frac{f_i(\mathbf{v})}{v_i} \right) + \frac{g_i(\mathbf{v})}{v_i} &= \frac{f_i(\mathbf{v})}{v_i} + \frac{g_i(\mathbf{v})}{v_i} + \frac{\zeta}{2} \\ &\leq -\frac{\zeta}{2}, \quad i = 1, \dots, n, \end{aligned}$$

where we have used (B.1) to get the inequality. For each i , it follows that

$$(B.3) \quad \gamma^{\frac{r_i+p}{r_{\max}}} \left(\frac{f_i(\mathbf{v})}{v_i} \right) + \frac{g_i(\mathbf{v})}{v_i} \leq -\frac{\zeta}{2},$$

where $\gamma = \max_{1 \leq i \leq n} \gamma_i$. Clearly, $\gamma \in (0, 1)$. The proof now proceeds in two steps:

1. First, we show that for any initial condition $\varphi(\cdot) \in \mathcal{C}([-\tau_{\max}, 0], \mathbb{R}_+^n)$, the corresponding solution $\mathbf{x}(t)$ of (3.1) satisfies $\mathbf{x}(t) \in S(0)$ for all $t \geq 0$, where the sets $S(m)$ are defined in (3.8).

2. By induction, we then prove that for each $m \in \mathbb{N}_0$, there exists a time $t_m \geq 0$ such that $\mathbf{x}(t)$ will enter the set $S(m)$ at t_m and remains in this set for all $t \geq t_m$.

Step 1. Since the homogeneous cooperative system (3.1) is positive, $x_i(t) \geq 0$ for all $i \in \{1, \dots, n\}$ and all $t \geq 0$. Let

$$(B.4) \quad z_i(t) = \left(\frac{x_i(t)}{v_i} \right)^{\frac{r_{\max}}{r_i}} - \|\varphi\|,$$

where $\|\varphi\|$ is defined in (3.9). From the definition of $\|\varphi\|$, $z_i(0) \leq 0$ for all i . We claim that $z_i(t) \leq 0$ for all $t \geq 0$. By contradiction, suppose this is not true. Then, there exist an index $j \in \{1, \dots, n\}$ and a time $t_1 \geq 0$ such that

$$(B.5) \quad \begin{aligned} z_i(t) &\leq 0, \quad i = 1, \dots, n, \quad t \in [0, t_1], \\ z_j(t_1) &= 0, \end{aligned}$$

and

$$(B.6) \quad D^+ z_j(t) \Big|_{t=t_1} \geq 0.$$

From (B.4) and (B.5), we have

$$\begin{aligned} x_i(t_1) &\leq (\lambda_\varphi)^{r_i} v_i, \quad i = 1, \dots, n, \quad i \neq j, \\ x_j(t_1) &= (\lambda_\varphi)^{r_j} v_j, \end{aligned}$$

where $\lambda_\varphi = \|\varphi\|^{\frac{1}{r_{\max}}}$. Since \mathbf{f} is cooperative and homogeneous of degree p with respect to the dilation map $\delta_\lambda^r(\mathbf{x})$, it follows from Proposition 2.2 that

$$(B.7) \quad f_j(\mathbf{x}(t_1)) \leq f_j(\delta_{\lambda_\varphi}^r(\mathbf{v})) = (\lambda_\varphi)^{r_j+p} f_j(\mathbf{v}).$$

If $t_1 - \tau(t_1) \in [0, t_1]$, then, from (B.5), we have $z_i(t_1 - \tau(t_1)) \leq 0$, which implies that $x_i(t_1 - \tau(t_1)) \leq (\lambda_\varphi)^{r_i} v_i$ for all i , or, equivalently,

$$\mathbf{x}(t_1 - \tau(t_1)) \leq \delta_{\lambda_\varphi}^r(\mathbf{v}).$$

Note also that if $t_1 - \tau(t_1) \in [-\tau_{\max}, 0]$, then $\mathbf{x}(t_1 - \tau(t_1)) = \varphi(t_1 - \tau(t_1))$, and hence, from the definition of $\|\varphi\|$, the above inequality still holds. As \mathbf{g} is nondecreasing and homogeneous, this in turn implies that

$$(B.8) \quad g_j(\mathbf{x}(t_1 - \tau(t_1))) \leq g_j(\delta_{\lambda_\varphi}^r(\mathbf{v})) = (\lambda_\varphi)^{r_j+p} g_j(\mathbf{v}).$$

The upper-right Dini-derivative of $z_j(t)$ along the trajectories of (3.1) at $t = t_1$ is given by

$$\begin{aligned} D^+ z_j(t) \Big|_{t=t_1} &= \left(\frac{r_{\max}}{r_j} \right) \left(\frac{x_j(t_1)}{v_j} \right)^{\left(\frac{r_{\max}}{r_j} - 1 \right)} \left(\frac{f_j(\mathbf{x}(t_1)) + g_j(\mathbf{x}(t_1 - \tau(t_1)))}{v_j} \right) \\ &\leq \left(\frac{r_{\max}}{r_j} \right) (\lambda_\varphi)^{r_{\max}-r_j} (\lambda_\varphi)^{r_j+p} \left(\frac{f_j(\mathbf{v}) + g_j(\mathbf{v})}{v_j} \right) \\ &= \left(\frac{r_{\max}}{r_j} \right) (\lambda_\varphi)^{r_{\max}+p} \left(\frac{f_j(\mathbf{v}) + g_j(\mathbf{v})}{v_j} \right), \end{aligned}$$

where we have used (B.7) and (B.8) to obtain the inequality. It follows from (3.10) that $D^+ z_j(t_1) < 0$, which contradicts (B.6). Therefore, $z_i(t) \leq 0$ for all i and all $t \geq 0$, and hence $V(\mathbf{x}(t)) \leq \|\varphi\|$ for $t \geq 0$. This shows that $\mathbf{x}(t) \in S(0)$ for all $t \geq 0$.

Step 2. According to the previous step, the induction hypothesis is true for $m = 0$. Now, assume that it holds for a given m , i.e., $V(\mathbf{x}(t)) \leq \gamma^m \|\varphi\|$ for all $t \geq t_m$. We will prove that there exists a finite time $t_{m+1} \geq 0$ such that $\mathbf{x}(t_{m+1}) \in S(m+1)$. By contradiction, suppose this is not true. Then,

$$(B.9) \quad \gamma^{m+1} \|\varphi\| \leq V(\mathbf{x}(t)) \leq \gamma^m \|\varphi\| \quad \forall t \geq t_m.$$

Let $i_t \in \{1, \dots, n\}$ be an index such that $V(\mathbf{x}(t)) = V_{i_t}(x_{i_t}(t))$, where

$$V_i(x_i) = \left(\frac{x_i}{v_i} \right)^{\frac{r_{\max}}{r_i}}.$$

The cooperativity and homogeneity of \mathbf{f} implies that

$$(B.10) \quad \begin{aligned} f_{i_t}(\mathbf{x}(t)) &\leq (V(\mathbf{x}(t)))^{\left(\frac{r_{i_t}+p}{r_{\max}} \right)} f_{i_t}(\mathbf{v}) \\ &\leq (\gamma^{m+1} \|\varphi\|)^{\left(\frac{r_{i_t}+p}{r_{\max}} \right)} f_{i_t}(\mathbf{v}) \quad \forall t \geq t_m, \end{aligned}$$

where the second inequality follows from (B.9) and the fact that $f_{i_t}(\mathbf{v}) < 0$. From Assumption 3.1, $\lim_{t \rightarrow \infty} t - \tau(t) = +\infty$. Thus, there exists sufficiently large $\hat{t}_m \geq t_m$ so that $t - \tau(t) \geq t_m$ for all $t \geq \hat{t}_m$. Since $x(t) \in S(m)$ for $t \geq t_m$, it follows that $x(t - \tau(t)) \in S(m)$ for all $t \geq \hat{t}_m$, implying that $V(\mathbf{x}(t - \tau(t))) \leq \gamma^m \|\varphi\|$ for $t \geq \hat{t}_m$, or, equivalently,

$$(B.11) \quad x_i(t - \tau(t)) \leq (\gamma^m \|\varphi\|)^{\left(\frac{r_i}{r_{\max}}\right)} v_i \quad \forall t \geq \hat{t}_m$$

for all i . As \mathbf{g} is nondecreasing and homogeneous, we then have

$$(B.12) \quad g_{i_t}(\mathbf{x}(t - \tau(t))) \leq (\gamma^m \|\varphi\|)^{\left(\frac{r_{i_t} + p}{r_{\max}}\right)} g_{i_t}(\mathbf{v}) \quad \forall t \geq \hat{t}_m.$$

Substituting (B.10) and (B.12) into the upper-right Dini-derivative of $V_{i_t}(x_{i_t})$ along the trajectories of (3.1) yields

$$(B.13) \quad \begin{aligned} & D^+ V_{i_t}(x_{i_t}) \\ &= \left(\frac{r_{\max}}{r_{i_t}}\right) \left(\frac{x_{i_t}(t)}{v_{i_t}}\right)^{\left(\frac{r_{\max}}{r_{i_t}} - 1\right)} \left(\frac{f_{i_t}(\mathbf{x}(t)) + g_{i_t}(\mathbf{x}(t - \tau(t)))}{v_{i_t}}\right) \\ &\leq \left(\frac{r_{\max}}{r_{i_t}}\right) \left(\frac{x_{i_t}(t)}{v_{i_t}}\right)^{\left(\frac{r_{\max}}{r_{i_t}} - 1\right)} (\gamma^m \|\varphi\|)^{\left(\frac{r_{i_t} + p}{r_{\max}}\right)} \left(\gamma^{\frac{r_{i_t} + p}{r_{\max}}} \left(\frac{f_{i_t}(\mathbf{v})}{v_{i_t}}\right) + \frac{g_{i_t}(\mathbf{v})}{v_{i_t}}\right) \\ &\leq -\left(\frac{r_{\max}}{r_{i_t}}\right) \left(\frac{x_{i_t}(t)}{v_{i_t}}\right)^{\left(\frac{r_{\max}}{r_{i_t}} - 1\right)} (\gamma^m \|\varphi\|)^{\left(\frac{r_{i_t} + p}{r_{\max}}\right)} \left(\frac{\zeta}{2}\right), \\ &\leq -\underbrace{\left(\frac{r_{\max}}{r_{i_t}}\right) (\gamma^{m+1} \|\varphi\|)^{\left(1 - \frac{r_{i_t}}{r_{\max}}\right)} (\gamma^m \|\varphi\|)^{\left(\frac{r_{i_t} + p}{r_{\max}}\right)}}_{\kappa} \left(\frac{\zeta}{2}\right) \quad \forall t \geq \hat{t}_m, \end{aligned}$$

where the last two inequalities follow from (B.3) and (B.9), respectively. Note that $\kappa > 0$. Since $V_i(x_i)$ is continuously differentiable on \mathbb{R} for each i , $V(\mathbf{x})$ is locally Lipschitz and

$$D^+ V(\mathbf{x}(t)) = \max_{j \in \mathcal{J}(t)} D^+ V_j(x_j(t)),$$

where $\mathcal{J}(t) = \{i \mid V_i(x_i(t)) = V(\mathbf{x}(t))\}$ [8]. It follows from (B.13) that

$$D^+ V(\mathbf{x}(t)) \leq -\kappa \quad \forall t \geq \hat{t}_m.$$

This together with (B.9) implies that

$$\begin{aligned} V(\mathbf{x}(t)) &\leq V(\mathbf{x}(\hat{t}_m)) - \kappa(t - \hat{t}_m) \\ &\leq \gamma^m \|\varphi\| - \kappa(t - \hat{t}_m) \quad \forall t \geq \hat{t}_m. \end{aligned}$$

It is immediate to see that the the right-hand side of the above inequality becomes smaller than $\gamma^{m+1} \|\varphi\|$ when

$$t \geq t_{m+1} = \hat{t}_m + \gamma^m \|\varphi\| \frac{1 - \gamma}{\kappa},$$

which contradicts (B.9). Thus, necessarily, $\mathbf{x}(t)$ reaches $S(m + 1)$ in a finite time.

We now prove that $\mathbf{x}(t)$ remains in $S(m+1)$ for all $t \geq t_{m+1}$. Let

$$(B.14) \quad w_i(t) = \left(\frac{x_i(t)}{v_i} \right)^{\frac{r_{\max}}{r_i}} - \gamma^{m+1} \|\varphi\|, \quad t \geq t_{m+1}.$$

Since $\mathbf{x}(t_{m+1}) \in S(m+1)$, $w_i(t_{m+1}) \leq 0$ for all i . We show that $w_i(t) \leq 0$ for all i and all $t \geq t_{m+1}$. If, by contradiction, this is not true, then there is an index $j \in \{1, \dots, n\}$ and a time $t_2 \geq t_{m+1}$ such that $w_i(t) \leq 0$ for $t \in [t_{m+1}, t_2]$, $w_j(t_2) = 0$, and

$$(B.15) \quad D^+ w_j(t) \Big|_{t=t_2} \geq 0.$$

From (B.14), we have

$$\begin{aligned} x_i(t_2) &\leq (\gamma^{m+1} \|\varphi\|)^{\frac{r_i}{r_{\max}}} v_i, \quad i = 1, \dots, n, \quad i \neq j, \\ x_j(t_2) &= (\gamma^{m+1} \|\varphi\|)^{\frac{r_j}{r_{\max}}} v_j. \end{aligned}$$

It now follows from cooperativity and homogeneity of \mathbf{f} that

$$(B.16) \quad f_j(\mathbf{x}(t_2)) \leq (\gamma^{m+1} \|\varphi\|)^{\frac{r_j+p}{r_{\max}}} f_j(\mathbf{v}).$$

Moreover, since $t_2 \geq t_{m+1} \geq \hat{t}_m$, it follows from (B.11) that

$$(B.17) \quad g_j(\mathbf{x}(t_2 - \tau(t_2))) \leq (\gamma^m \|\varphi\|)^{\frac{r_j+p}{r_{\max}}} g_j(\mathbf{v}),$$

where we have used the fact that \mathbf{g} is nondecreasing and homogeneous. The upper-right Dini-derivative of $w_j(t)$ along the trajectories of (3.1) at $t = t_2$ is given by

$$\begin{aligned} &D^+ w_j(t) \Big|_{t=t_2} \\ &\leq \left(\frac{r_{\max}}{r_j} \right) \left(\frac{x_j(t_2)}{v_j} \right)^{\left(\frac{r_{\max}}{r_j} - 1 \right)} \left(\gamma^m \|\varphi\| \right)^{\frac{r_j+p}{r_{\max}}} \left(\gamma^{\frac{r_j+p}{r_{\max}}} \left(\frac{f_j(\mathbf{v})}{v_j} \right) + \frac{g_j(\mathbf{v})}{v_j} \right) \\ &< 0, \end{aligned}$$

where we have used (B.16) and (B.17) to get the first inequality and (B.3) to obtain the second inequality. This contradicts (B.15), and hence $w_i(t) \leq 0$ for all i and all $t \geq t_{m+1}$. It follows that $V(\mathbf{x}(t)) \leq \gamma^{m+1} \|\varphi\|$ for $t \geq t_{m+1}$, or, equivalently, $\mathbf{x}(t) \in S(m+1)$ for all $t \geq t_{m+1}$.

In summary, we conclude that for each $m \in \mathbb{N}_0$, there exists $t_m \geq 0$ such that $\mathbf{x}(t) \in S(m)$ for all $t \geq t_m$. Since $\gamma < 1$, γ^m approaches zero as $m \rightarrow \infty$. Therefore, the origin is globally asymptotically stable.

(ii) \Rightarrow (iii). Assume that (3.1) is globally asymptotically stable for all delays satisfying Assumption 3.1. Particularly, let $\tau(t) = 0$. Then, $\dot{\mathbf{x}}(t) = \mathbf{f}(\mathbf{x}(t)) + \mathbf{g}(\mathbf{x}(t))$ is asymptotically stable.

(iii) \Rightarrow (i). As $\mathbf{f} + \mathbf{g}$ is a cooperative vector field, it follows from [39, Proposition 3.10, Theorem 3.12] that there is some vector $\mathbf{v} > \mathbf{0}$ satisfying (3.10).

Appendix C. Proof of Theorem 3.5. Let $\mathbf{v} > \mathbf{0}$ be a vector satisfying (3.11). According to Theorem 3.3, the homogeneous cooperative system (3.1) is globally

asymptotically stable for all nonnegative initial conditions and for all delays satisfying Assumption 3.1. We will prove that it is also globally μ -stable. From Remark 3.1, there exists a constant $T_0 > 0$ large enough such that

$$(C.1) \quad t - \tau(t) > 0 \quad \forall t > T_0.$$

By condition (iv), we can find a sufficiently large constant $T_1 > 0$ such that for all $t > T_1$ and all $i \in \{1, \dots, n\}$,

$$\left(\frac{r_{\max}}{r_i}\right) \left(\left(\frac{f_i(\mathbf{v})}{v_i}\right) + \left(\frac{\mu(t)}{\mu(t-\tau(t))}\right)^{\frac{r_i+p}{r_{\max}}} \left(\frac{g_i(\mathbf{v})}{v_i}\right) \right) + \frac{\dot{\mu}(t)}{(\mu(t))^{1-\frac{p}{r_{\max}}}} < 0.$$

Since $\mu(t)$ is positive and nondecreasing on \mathbb{R}_+ , it follows that

$$(C.2) \quad \epsilon \left(\frac{r_{\max}}{r_i}\right) \left(\left(\frac{f_i(\mathbf{v})}{v_i}\right) + \left(\frac{\mu(t)}{\mu(t-\tau(t))}\right)^{\frac{r_i+p}{r_{\max}}} \left(\frac{g_i(\mathbf{v})}{v_i}\right) \right) + \frac{\dot{\mu}(t)}{(\mu(t))^{1-\frac{p}{r_{\max}}}} < 0$$

holds for any $\epsilon \geq 1$. Let $M = \max\{1, \mu(\bar{T})\|\varphi\|\}$, where $\bar{T} = \max\{T_0, T_1\} + 1$, and $\|\varphi\|$ is defined in (3.9). According to Theorem 3.3, $V(\mathbf{x}(t)) \leq \|\varphi\|$ for all $t \geq 0$. Thus,

$$(C.3) \quad \begin{aligned} \sup_{0 \leq t \leq \bar{T}} \{\mu(t)V(\mathbf{x}(t))\} &\leq \sup_{0 \leq t \leq \bar{T}} \{\mu(t)\|\varphi\|\} \\ &= \mu(\bar{T})\|\varphi\| \\ &\leq M, \end{aligned}$$

where we have used condition (ii) to get the equality. It follows that

$$(C.4) \quad \mu(t)V(\mathbf{x}(t)) \leq M \quad \forall t \in [0, \bar{T}].$$

In order to prove global μ -stability, we will show that (C.4) also holds for all $t \geq \bar{T}$. By contradiction, suppose this is not true. Then, there exist an index $j \in \{1, \dots, n\}$ and a time $t_1 \geq \bar{T}$ such that

$$(C.5) \quad \mu(t)V(\mathbf{x}(t)) \leq M, \quad t \in [0, t_1],$$

$$(C.6) \quad \mu(t_1) \left(\frac{x_j(t_1)}{v_j}\right)^{\frac{r_{\max}}{r_j}} = M,$$

$$(C.7) \quad D^+ \mu(t) \left(\frac{x_j(t)}{v_j}\right)^{\frac{r_{\max}}{r_j}} \Big|_{t=t_1} \geq 0.$$

From (C.5) and (C.6), we have

$$\begin{aligned} x_i(t_1) &\leq \left(\frac{M}{\mu(t_1)}\right)^{\frac{r_i}{r_{\max}}} v_i, \quad i = 1, \dots, n, \quad i \neq j, \\ x_j(t_1) &= \left(\frac{M}{\mu(t_1)}\right)^{\frac{r_j}{r_{\max}}} v_j. \end{aligned}$$

Now, as \mathbf{f} is cooperative and homogeneous, it follows from Proposition 2.2 that

$$(C.8) \quad f_j(\mathbf{x}(t_1)) \leq \left(\frac{M}{\mu(t_1)}\right)^{\frac{r_j+p}{r_{\max}}} f_j(\mathbf{v}).$$

Since $t_1 \geq \bar{T} > T_0$, it follows from (C.1) that $t_1 \geq t_1 - \tau(t_1) > 0$. Hence, from (C.5), we have

$$\mu(t_1 - \tau(t_1))V(\mathbf{x}(t_1 - \tau(t_1))) \leq M.$$

As \mathbf{g} is nondecreasing and homogeneous, this in turn implies

$$(C.9) \quad g_j(\mathbf{x}(t_1 - \tau(t_1))) \leq \left(\frac{M}{\mu(t_1 - \tau(t_1))} \right)^{\frac{r_j+p}{r_{\max}}} g_j(\mathbf{v}).$$

We then have

$$\begin{aligned} D^+ \mu(t) \left(\frac{x_j(t)}{v_j} \right)^{\frac{r_{\max}}{r_j}} \Big|_{t=t_1} &= \mu(t_1) \left(\frac{r_{\max}}{r_j} \right) \left(\frac{x_j(t_1)}{v_j} \right)^{\left(\frac{r_{\max}}{r_j} - 1 \right)} \frac{\dot{x}_j(t_1)}{v_j} + \dot{\mu}(t_1) \left(\frac{x_j(t_1)}{v_j} \right)^{\frac{r_{\max}}{r_j}} \\ &= \mu(t_1) \left(\frac{r_{\max}}{r_j} \right) \left(\frac{M}{\mu(t_1)} \right)^{\left(1 - \frac{r_j}{r_{\max}} \right)} \left(\frac{f_j(\mathbf{x}(t_1)) + g_j(\mathbf{x}(t_1 - \tau(t_1)))}{v_j} \right) + M \frac{\dot{\mu}(t_1)}{\mu(t_1)} \\ &\leq \frac{M}{\left(\mu(t_1) \right)^{\frac{p}{r_{\max}}}} \\ &\quad \times \left\{ M^{\frac{p}{r_{\max}}} \left(\frac{r_{\max}}{r_j} \right) \left(\left(\frac{f_j(\mathbf{v})}{v_j} \right) + \left(\frac{\mu(t_1)}{\mu(t_1 - \tau(t_1))} \right)^{\frac{r_j+p}{r_{\max}}} \left(\frac{g_j(\mathbf{v})}{v_j} \right) \right) + \frac{\dot{\mu}(t_1)}{\mu(t_1)^{1-\frac{p}{r_{\max}}}} \right\}, \end{aligned}$$

where we have used (C.6) to get the second equality, and (C.8)–(C.9) to obtain the inequality. Since $M \geq 1$ and $t_1 \geq \bar{T} > T_1$, it now follows from (C.2) that

$$D^+ \mu(t) \left(\frac{x_j(t)}{v_j} \right)^{\frac{r_{\max}}{r_j}} \Big|_{t=t_1} < 0,$$

which contradicts (C.7). We conclude that $\mu(t)V(\mathbf{x}(t)) \leq M$ for all $t \geq \bar{T}$, and hence

$$V(\mathbf{x}(t)) \leq \frac{M}{\mu(t)}, \quad t \geq 0.$$

This completes the proof.

Appendix D. Proof of Corollary 3.6. (i) Assume that $p = 0$. According to Remark 3.3, (3.14) has a unique positive solution η_i . Pick a constant η satisfying $0 < \eta < \min_{1 \leq i \leq n} \eta_i$. Since the left-hand side of (3.14) is strictly monotonically increasing in $\eta_i > 0$, we have

$$(D.1) \quad \left(\frac{r_{\max}}{r_i} \right) \left(\left(\frac{f_i(\mathbf{v})}{v_i} \right) + \left(e^{\eta r_{\max}} \right)^{\frac{r_i}{r_{\max}}} \left(\frac{g_i(\mathbf{v})}{v_i} \right) \right) + \eta < 0, \quad i = 1, \dots, n.$$

Now, let $\mu(t) = e^{\eta t}$. One can verify that $\mu(t)$ satisfies conditions (i)–(iii) of Theorem 3.5. Moreover,

$$\lim_{t \rightarrow \infty} \frac{\dot{\mu}(t)}{\mu(t)} = \eta,$$

and

$$\lim_{t \rightarrow \infty} \frac{\mu(t)}{\mu(t - \tau(t))} \leq \lim_{t \rightarrow \infty} \frac{e^{\eta t}}{e^{\eta(t - \tau_{\max})}} = e^{\eta \tau_{\max}},$$

where the inequality holds since $\tau(t) \leq \tau_{\max}$ and $\mu(t)$ is nondecreasing. It follows from (D.1) and the above observations that condition (iv) of Theorem 3.5 is also satisfied. Hence, the solution $\mathbf{x}(t)$ of (3.1) satisfies (3.13).

(ii) If $p > 0$, we can pick $\mu(t) = (\theta t + 1)^{\frac{r_{\max}}{p}}$. The rest of the proof is similar to the one for $p = 0$ and thus omitted.

Appendix E. Proof of Corollary 3.7. (i) Assume that $p = 0$. The left-hand side of (3.17) is strictly monotonically increasing in $\xi_i > 0$, which implies that

$$\left(\frac{f_i(\mathbf{v})}{v_i} \right) + \left(\frac{1}{1 - \alpha} \right)^{\frac{r_i}{r_{\max}} \xi} \left(\frac{g_i(\mathbf{v})}{v_i} \right) < 0, \quad i = 1, \dots, n,$$

where $\xi \in (0, \min_{1 \leq i \leq n} \xi_i)$. Now, letting $\mu(t) = t^\xi$, the rest of the proof is similar to the one of Corollary 3.6 and thus omitted.

(ii) For $p > 0$, we can choose $\mu(t) = t^{\frac{r_{\max}}{p} \beta}$, where β satisfies (3.18).

Appendix F. Proof of Theorem 4.3. (i) \Rightarrow (ii). Let $\mathbf{v} > \mathbf{0}$ be a vector such that (4.3) holds and let

$$(F.1) \quad \gamma = \max_{1 \leq i \leq n} \left(\frac{f_i(\mathbf{v}) + g_i(\mathbf{v})}{v_i} \right)^{\frac{r_{\max}}{r_i}}.$$

Note that $0 \leq \gamma < 1$. The proof will be broken up into two steps:

1. We show that for any nonnegative initial condition $\varphi(\cdot)$, $\mathbf{x}(k) \in S(0)$ for all $k \in \mathbb{N}_0$, where the sets $S(m)$ are defined in (3.8).

2. We then use induction to show that for each $m \in \mathbb{N}_0$, there exists $k_m \in \mathbb{N}_0$ such that $\mathbf{x}(k) \in S(m)$ for all $k \geq k_m$.

Step 1. Since the initial state $\mathbf{x}(0)$ satisfies $V(\mathbf{x}(0)) \leq \|\varphi\|$, $\mathbf{x}(0) \in S(0)$. Assume for induction that $\mathbf{x}(k) \in S(0)$ holds for all k up to some $k \in \mathbb{N}_0$. Thus,

$$x_i(k) \leq \|\varphi\|^{\frac{r_i}{r_{\max}}} v_i, \quad k \in \{-d_{\max}, \dots, \bar{k}\},$$

for all i . As \mathbf{f} and \mathbf{g} are homogeneous of degree zero and nondecreasing on \mathbb{R}_+^n , it follows that

$$(F.2) \quad \begin{aligned} f_i(\mathbf{x}(\bar{k})) &\leq \|\varphi\|^{\frac{r_i}{r_{\max}}} f_i(\mathbf{v}), \\ g_i(\mathbf{x}(\bar{k} - d(\bar{k}))) &\leq \|\varphi\|^{\frac{r_i}{r_{\max}}} g_i(\mathbf{v}). \end{aligned}$$

For each $i \in \{1, \dots, n\}$, we then have

$$\begin{aligned} \frac{1}{v_i} x_i(\bar{k} + 1) &= \frac{f_i(\mathbf{x}(\bar{k})) + g_i(\mathbf{x}(\bar{k} - d(\bar{k})))}{v_i} \\ &\leq \|\varphi\|^{\frac{r_i}{r_{\max}}} \left(\frac{f_i(\mathbf{v}) + g_i(\mathbf{v})}{v_i} \right) \\ &\leq \|\varphi\|^{\frac{r_i}{r_{\max}}}, \end{aligned}$$

where we have used (F.2) to get the first inequality and (4.3) to obtain the second inequality. It follows that $V(\mathbf{x}(\bar{k} + 1)) \leq \|\boldsymbol{\varphi}\|$, which completes the induction proof. Therefore, $\mathbf{x}(k) \in S(0)$ for all $k \in \mathbb{N}_0$.

Step 2. Assume for induction that $\mathbf{x}(k) \in S(m)$ for all $k \geq k_m$, i.e.,

$$x_i(k) \leq (\gamma^m \|\boldsymbol{\varphi}\|)^{\frac{r_i}{r_{\max}}} v_i, \quad k \geq k_m.$$

We will show that there exists $k_{m+1} \in \mathbb{N}_0$ such that $\mathbf{x}(k) \in S(m+1)$ for all $k \geq k_{m+1}$. Since \mathbf{f} is homogeneous of degree zero and nondecreasing on \mathbb{R}_+^n , we have

$$(F.3) \quad f_i(\mathbf{x}(k)) \leq (\gamma^m \|\boldsymbol{\varphi}\|)^{\frac{r_i}{r_{\max}}} f_i(\mathbf{v}), \quad k \geq k_m.$$

According to Assumption 4.1, $\lim_{k \rightarrow \infty} k - d(k) = +\infty$, so one can find a sufficiently large $\bar{k}_m \geq k_m$, $\bar{k}_m \in \mathbb{N}$, such that $k - d(k) \geq k_m$ for all $k \geq \bar{k}_m$. Moreover, since $\mathbf{x}(k) \in S(m)$ for $k \geq k_m$, we have $\mathbf{x}(k - d(k)) \in S(m)$ for all $k \geq \bar{k}_m$, which implies that $V(\mathbf{x}(k - d(k))) \leq \gamma^m \|\boldsymbol{\varphi}\|$, or, equivalently,

$$x_i(k - d(k)) \leq (\gamma^m \|\boldsymbol{\varphi}\|)^{\frac{r_i}{r_{\max}}} v_i, \quad k \geq \bar{k}_m.$$

As \mathbf{g} is homogeneous of degree zero and nondecreasing on \mathbb{R}_+^n , it follows that

$$(F.4) \quad g_i(\mathbf{x}(k - d(k))) \leq (\gamma^m \|\boldsymbol{\varphi}\|)^{\frac{r_i}{r_{\max}}} g_i(\mathbf{v}), \quad k \geq \bar{k}_m.$$

For each $i \in \{1, \dots, n\}$, we then have

$$\begin{aligned} \frac{1}{v_i} x_i(k+1) &\leq (\gamma^m \|\boldsymbol{\varphi}\|)^{\frac{r_i}{r_{\max}}} \left(\frac{f_i(\mathbf{v}) + g_i(\mathbf{v})}{v_i} \right) \\ &\leq (\gamma^{m+1} \|\boldsymbol{\varphi}\|)^{\frac{r_i}{r_{\max}}} \quad \forall k \geq \bar{k}_m, \end{aligned}$$

where we have used (F.3) and (F.4) to get the first inequality, and (F.1) to obtain the second inequality. Therefore,

$$V(\mathbf{x}(k+1)) \leq \gamma^{m+1} \|\boldsymbol{\varphi}\| \quad \forall k \geq \bar{k}_m,$$

which implies that $\mathbf{x}(k+1) \in S(m+1)$ for all $k \geq \bar{k}_m$. Thus,

$$\mathbf{x}(k) \in S(m+1) \quad \forall k \geq \bar{k}_m + 1.$$

Letting $k_{m+1} = \bar{k}_m + 1$, the induction proof is complete.

In summary, we conclude that for each m , there exists k_m such that $\mathbf{x}(k) \in S(m)$ for all $k \geq k_m$. Since $\gamma < 1$, γ^m approaches zero as $m \rightarrow +\infty$. Hence, the homogeneous nondecreasing system (4.1) is globally asymptotically stable.

(ii) \Rightarrow (iii). Suppose that (4.1) is asymptotically stable for all delays satisfying Assumption 4.1. Particularly, let $d(k) = 0$. Then, $\mathbf{x}(k+1) = \mathbf{f}(\mathbf{x}(k)) + \mathbf{g}(\mathbf{x}(k))$ is asymptotically stable.

(iii) \Rightarrow (i). Since $\mathbf{f} + \mathbf{g}$ is continuous, nondecreasing on \mathbb{R}_+^n and $(\mathbf{f} + \mathbf{g})(\mathbf{0}) = \mathbf{0}$, the conclusion follows from [9, Propositions 5.2 and 5.6].

Appendix G. Proof of Corollary 4.4. Note that since $\boldsymbol{\varphi}(k) \leq \mathbf{v}$ for all $k \in \{-d_{\max}, \dots, 0\}$, we have $\|\boldsymbol{\varphi}\| \leq 1$. The proof is similar to the one of Theorem 4.3 and thus omitted.

Appendix H. Proof of Theorem 4.6. Let $\mathbf{v} > \mathbf{0}$ be a vector satisfying (4.4). According to Theorem 4.3, the homogeneous nondecreasing system (4.1) with time

delays satisfying Assumption 4.1 is globally asymptotically stable. We will prove that it is also globally μ -stable. From Remark 4.1, there exists $T_0 \in \mathbb{N}$ such that

$$(H.1) \quad k - d(k) > 0 \quad \forall k > T_0.$$

From condition (iv), one can find a sufficiently large constant $T_1 \in \mathbb{N}$, such that for all $k > T_1$, we have

$$(H.2) \quad \left(\frac{\mu(k+1)}{\mu(k)}\right)^{\frac{r_i}{r_{\max}}} \left(\frac{f_i(\mathbf{v})}{v_i}\right) + \left(\frac{\mu(k+1)}{\mu(k-d(k))}\right)^{\frac{r_i}{r_{\max}}} \left(\frac{g_i(\mathbf{v})}{v_i}\right) < 1.$$

Let $M = \mu(\bar{T})\|\varphi\|$, where $\bar{T} = \max\{T_0, T_1\} + 1$, and $\|\varphi\|$ is defined in (3.9). We now use induction to prove that

$$(H.3) \quad V(\mathbf{x}(k)) \leq \frac{M}{\mu(k)} \quad \forall k \in \mathbb{N}.$$

According to Theorem 4.3, $V(\mathbf{x}(k)) \leq \|\varphi\|$ for all $k \in \mathbb{N}$. Thus,

$$(H.4) \quad \begin{aligned} \max_{1 \leq k \leq \bar{T}} \{\mu(k)V(\mathbf{x}(k))\} &\leq \max_{1 \leq k \leq \bar{T}} \{\mu(k)\}\|\varphi\| \\ &= \mu(\bar{T})\|\varphi\| \\ &= M, \end{aligned}$$

where we have used condition (ii) to get the first equality. It follows from (H.4) that (H.3) is true for $k \in \{1, \dots, \bar{T}\}$. Next, assume for induction that (H.3) holds for all k up to some \bar{k} , where $\bar{k} \geq \bar{T}$. Thus,

$$0 \leq \left(\frac{x_i(k)}{v_i}\right)^{\frac{r_{\max}}{r_i}} \leq \frac{M}{\mu(k)}, \quad k = 1, \dots, \bar{k},$$

which implies that

$$(H.5) \quad 0 \leq x_i(\bar{k}) \leq \left(\frac{M}{\mu(\bar{k})}\right)^{\frac{r_i}{r_{\max}}} v_i.$$

Since $\bar{k} \geq \bar{T} > T_0$, it follows from (H.1) that $\bar{k} - d(\bar{k}) \in \{1, \dots, \bar{k}\}$. Hence,

$$(H.6) \quad 0 \leq x_i(\bar{k} - d(\bar{k})) \leq \left(\frac{M}{\mu(\bar{k} - d(\bar{k}))}\right)^{\frac{r_i}{r_{\max}}} v_i.$$

As \mathbf{f} and \mathbf{g} are homogeneous of degree zero and nondecreasing on \mathbb{R}_+^n , it follows from (H.5) and (H.6) that

$$(H.7) \quad \begin{aligned} f_i(\mathbf{x}(\bar{k})) &\leq \left(\frac{M}{\mu(\bar{k})}\right)^{\frac{r_i}{r_{\max}}} f_i(\mathbf{v}), \\ g_i(\mathbf{x}(\bar{k} - d(\bar{k}))) &\leq \left(\frac{M}{\mu(\bar{k} - d(\bar{k}))}\right)^{\frac{r_i}{r_{\max}}} g_i(\mathbf{v}). \end{aligned}$$

We now show that $\mathbf{x}(\bar{k} + 1)$ satisfies (H.3). For each $i \in \{1, \dots, n\}$,

$$\begin{aligned} \frac{1}{v_i} x_i(\bar{k} + 1) &= \frac{f_i(\mathbf{x}(\bar{k})) + g_i(\mathbf{x}(\bar{k}) - d(\bar{k}))}{v_i} \\ &\leq \left(\frac{M}{\mu(\bar{k})} \right)^{\frac{r_i}{r_{\max}}} \left(\frac{f_i(\mathbf{v})}{v_i} \right) + \left(\frac{M}{\mu(\bar{k}) - d(\bar{k})} \right)^{\frac{r_i}{r_{\max}}} \left(\frac{g_i(\mathbf{v})}{v_i} \right) \\ &\leq \left(\frac{M}{\mu(\bar{k} + 1)} \right)^{\frac{r_i}{r_{\max}}}, \end{aligned}$$

where we have used (H.7) to get the first inequality and (H.2) to obtain the second inequality. Therefore,

$$V(\mathbf{x}(\bar{k} + 1)) \leq \frac{M}{\mu(\bar{k} + 1)},$$

and the induction proof is complete.

Acknowledgment. The authors would like to thank the anonymous associate editor and reviewers for suggestions that helped to improve the quality of the paper.

REFERENCES

- [1] D. AEYELS AND P. D. LEENHEER, *Extension of the Perron–Frobenius theorem to homogeneous systems*, SIAM J. Control Optim., 41 (2002), pp. 563–582.
- [2] A. BERMAN AND R. J. PLEMMONS, *Nonnegative Matrices in the Mathematical Sciences*, Academic Press, New York, 1979.
- [3] H. BOCHE AND M. SCHUBERT, *The structure of general interference functions and applications*, IEEE Trans. Inform. Theory, 54 (2008), pp. 4980–4990.
- [4] V. BOKHARAE, O. MASON, AND M. VERWOERD, *D-stability and delay-independent stability of homogeneous cooperative systems*, IEEE Trans. Automat. Control, 55 (2010), pp. 2882–2885.
- [5] C. BRIAT, *Robust stability and stabilization of uncertain linear positive systems via integral linear constraints: L1-gain and L∞-gain characterization*, Internat. J. Robust Nonlinear Control, 23 (2013), pp. 1932–1954.
- [6] T. CHARALAMBOUS, I. LESTAS, AND G. VINNICOMBE, *On the stability of the Foschini–Miljanic algorithm with time-delays*, in Proceedings of the 47th IEEE Conference on Decision and Control, 2008, pp. 2991–2996.
- [7] T. CHEN AND L. WANG, *Global μ-stability of delayed neural networks with unbounded time-varying delays*, IEEE Trans. Neural Networks, 18 (2007), pp. 1836–1840.
- [8] J. M. DANSKIN, *The theory of max–min, with applications*, SIAM J. Appl. Math., 14 (1966), pp. 641–664.
- [9] S. DASHKOVSKIY, B. RÜFFER, AND F. WIRTH, *Discrete-time monotone systems: Criteria for global asymptotic stability and applications*, in Proceedings of the 17th International Symposium on Mathematical Theory of Networks and Systems, 2006, pp. 89–97.
- [10] L. FAINSHIL, M. MARGALOT, AND P. CHIGANSKY, *On the stability of positive linear switched systems under arbitrary switching laws*, IEEE Trans. Automat. Control, 54 (2009), pp. 897–899.
- [11] L. FARINA AND S. RINALDI, *Positive Linear Systems: Theory and Applications*, Wiley, New York, 2000.
- [12] H. R. FEYZMAHDAVIAN, T. CHARALAMBOUS, AND M. JOHANSSON, *Asymptotic stability and decay rates of positive linear systems with unbounded delays*, in Proceedings of the 52th IEEE Conference on Decision and Control, 2013, pp. 6337–6342.
- [13] H. R. FEYZMAHDAVIAN, T. CHARALAMBOUS, AND M. JOHANSSON, *Exponential stability of homogeneous positive systems of degree one with time-varying delays*, IEEE Trans. Automat. Control, 59 (2014), pp. 1594–1599.
- [14] E. FORNASINI AND M. E. VALCHER, *Reachability of a class of discrete-time positive switched systems*, SIAM J. Control Optim., 49 (2011), pp. 162–184.

- [15] E. FRIDMAN AND U. SHAKED, *An improved stabilization method for linear time-delay systems*, IEEE Trans. Automat. Control, 47 (2002), pp. 1931–1937.
- [16] H. GAO AND T. CHEN, *New results on stability of discrete-time systems with time-varying state delay*, IEEE Trans. Automat. Control, 52 (2007), pp. 328–334.
- [17] W. M. HADDAD AND V. CHELLABOINA, *Stability theory for non-negative and compartmental dynamical systems with time delay*, Systems Control Lett., 51 (2004), pp. 355–361.
- [18] W. M. HADDAD, V. CHELLABOINA, AND Q. HUI, *Nonnegative and Compartmental Dynamical Systems*, Princeton University Press, Princeton, NJ, 2010.
- [19] J. K. HALE AND S. M. V. LUNEL, *Introduction to Functional Differential Equations*, Springer-Verlag, Berlin, 1993.
- [20] Y. HE, Q. G. WANG, C. LIN, AND M. WU, *Delay-range-dependent stability for systems with time-varying delay*, Automatica J. IFAC, 43 (2007), pp. 371–376.
- [21] Y. HINO, S. MURAKAMI, AND T. NAITO, *Functional Differential Equations with Infinite Delay*, Springer-Verlag, Berlin, 1991.
- [22] J. A. JACQUEZ AND C. P. SIMON, *Qualitative theory of compartmental systems*, SIAM Rev., 35 (1993), pp. 43–79.
- [23] L. F. KNORN, O. MASON, AND R. SHORTEN, *On linear co-positive Lyapunov functions for sets of linear positive systems*, Automatica J. IFAC, 45 (2009), pp. 1943–1947.
- [24] P. D. LEENHEER AND D. AEYELS, *Stabilization of positive linear systems*, Systems Control Lett., 44 (2001), pp. 259–271.
- [25] P. LI, J. LAM, Z. WANG, AND P. DATE, *Positivity-preserving H_∞ model reduction for positive systems*, Automatica J. IFAC, 47 (2011), pp. 1504–1511.
- [26] X. LIU AND C. DANG, *Stability analysis of positive switched linear systems with delays*, IEEE Trans. Automat. Control, 56 (2011), pp. 1684–1690.
- [27] X. LIU AND J. LAM, *Relationships between asymptotic stability and exponential stability of positive delay systems*, Int. J. Gen. Syst., 42 (2013), pp. 224–238.
- [28] X. LIU, W. YU, AND L. WANG, *Stability analysis of positive systems with bounded time-varying delays*, IEEE Trans. Circuits Syst. II, 56 (2009), pp. 600–604.
- [29] X. LIU, W. YU, AND L. WANG, *Stability analysis for continuous-time positive systems with time-varying delays*, IEEE Trans. Automat. Control, 55 (2010), pp. 1024–1028.
- [30] O. MASON AND R. SHORTEN, *On linear copositive Lyapunov functions and the stability of switched positive linear systems*, IEEE Trans. Automat. Control, 52 (2007), pp. 1346–1349.
- [31] O. MASON AND M. VERWOERD, *Observations on the stability of cooperative systems*, Systems Control Lett., 58 (2009), pp. 461–467.
- [32] P. H. A. NGOC, *A Perron–Frobenius theorem for a class of positive quasi-polynomial matrices*, Appl. Math. Lett., 19 (2006), pp. 747–751.
- [33] P. H. A. NGOC, *Stability of positive differential systems with delay*, IEEE Trans. Automat. Control, 58 (2013), pp. 203–209.
- [34] P. H. A. NGOC, S. MURAKAMI, T. NAITO, J. S. SHIN, AND Y. NAGABUCHI, *On positive linear Volterra–Stieltjes differential systems*, Integral Equations Operator Theory, 64 (2009), pp. 325–355.
- [35] M. M. PEET, A. PAPACHRISTODOULOU, AND S. LALL, *Positive forms and stability of linear time-delay systems*, SIAM J. Control Optim., 47 (2009), pp. 3237–3258.
- [36] M. A. RAMI, *Stability analysis and synthesis for linear positive systems with time-varying delays*, in Proceedings of the 3rd International Symposium on Positive Systems, 2009, pp. 205–216.
- [37] M. A. RAMI AND F. TADEO, *Controller synthesis for positive linear systems with bounded controls*, IEEE Trans. Circuits Syst. II, 54 (2007), pp. 151–155.
- [38] A. RANTZER, *Distributed control of positive systems*, in Proceedings of the 50th IEEE Conference on Decision and Control, 2011, pp. 6608–6611.
- [39] B. S. RÜFFER, C. M. KELLETT, AND S. R. WELLER, *Connection between cooperative positive systems and integral input-to-state stability of large-scale systems*, Automatica J. IFAC, 46 (2010), pp. 1019–1027.
- [40] R. SIPAHI, S.-I. NICULESCU, C. T. ABDALLAH, W. MICHIELS, AND K. GU, *Stability and stabilization of systems with time delay*, IEEE Control Syst. Mag., 31 (2011), pp. 38–65.
- [41] H. SMITH, *Monotone Dynamical Systems: An Introduction to the Theory of Competitive and Cooperative Systems*, AMS, Providence, RI, 1995.
- [42] Y. SUN, *Delay-independent stability of switched linear systems with unbounded time-varying delays*, Abstr. Appl. Anal., 2012 (2012), pp. 1–11.
- [43] T. TANAKA AND C. LANGBORT, *The bounded real lemma for internally positive systems and H -infinity structured static state feedback*, IEEE Trans. Automat. Control, 56 (2011), pp. 2218–2223.

- [44] M. E. VALCHER, *On the internal stability and asymptotic behavior of 2-D positive systems*, IEEE Trans. Circuits Syst. I, 44 (1997), pp. 602–613.
- [45] J. M. VAN DEN HOF, *Positive linear observers for linear compartmental systems*, SIAM J. Control Optim., 36 (1998), pp. 590–608.
- [46] M. WU, Y. HE, J. H. SHE, AND G. P. LIU, *Delay-dependent criteria for robust stability of time-varying delay systems*, Automatica J. IFAC, 40 (2004), pp. 1435–1439.
- [47] A. ZAPPAVIGNA, T. CHARALAMBOUS, AND F. KNORN, *Unconditional stability of the Foschini–Miljanic algorithm*, Automatica J. IFAC, 48 (2012), pp. 219–224.